

## LABELED OLIGONUCLEOTIDES, METHODS FOR MAKING SAME, AND COMPOUNDS USEFUL THEREFOR

### FIELD OF THE INVENTION

The present invention is directed to labeled oligonucleotides, methods for making the same,  
10 and compounds useful therefor. More specifically, this invention relates to oligonucleotides  
selectively functionalized at one or more of the 3'-terminal nucleotide, 5'-terminal nucleotide, and  
internucleotides with conjugate groups, methods for making the same, and compounds useful  
therefor.

### 15 BACKGROUND OF THE INVENTION

Oligonucleotides and their analogs have been developed and used in molecular biology in  
a variety of procedures as probes, primers, linkers, adapters, and gene fragments. The  
widespread use of such oligonucleotides has increased the demand for rapid, inexpensive and  
efficient procedures for their modification and synthesis. Early synthetic approaches to  
20 oligonucleotide synthesis included phosphodiester and phosphotriester chemistries. Khorana et  
al., *J. Molec. Biol.* 72, 209, 1972; Reese, *Tetrahedron Lett.* 34, 3143-3179, 1978. These  
approaches eventually gave way to more efficient modern methods, such as the use of  
phosphoramidites and H-phosphonates. Beaucage and Caruthers, *Tetrahedron Lett.*, 22, 1859-  
1862, 1981; Agrawal and Zamecnik, U.S. Patent No. 5,149,798, issued 1992.

25 The chemical literature discloses numerous processes for coupling nucleosides through  
phosphorous-containing covalent linkages to produce oligonucleotides of defined sequence. One  
of the most popular processes is the phosphoramidite technique (*see, e.g., Advances in the  
Synthesis of Oligonucleotides by the Phosphoramidite Approach*, Beaucage, S.L.; Iyer, R.P.,  
*Tetrahedron*, 1992, 48, 2223-2311 and references cited therein), wherein a nucleoside or  
30 oligonucleotide having a free hydroxyl group is reacted with a protected cyanoethyl  
phosphoramidite monomer in the presence of a weak acid to form a phosphite-linked structure.  
Oxidation of the phosphite linkage followed by hydrolysis of the cyanoethyl group yields the  
desired phosphodiester or phosphorothioate linkage.

The phosphoramidite technique, however, has significant disadvantages. For example,  
35 cyanoethyl phosphoramidite monomers are quite expensive. Although considerable quantities of

monomer go unreacted in a typical phosphoramidite coupling, unreacted monomer can be recovered, if at all, only with great difficulty.

The ability of the acylaminoethyl group to serve as a protecting group for certain phosphate diesters was first observed by Ziodrou and Schmir. Zioudrou et al., *J. Amer. Chem. Soc.*, 85, 3258, 1963. A version of this method was extended to the solid phase synthesis of oligonucleotide dimers, and oligomers with oxaphospholidine nucleoside building blocks as substitutes for conventional phosphoramidites. Iyer et al., *Tetrahedron Lett.*, 39, 2491-2494, 1998; PCT International Publication WO/9639413, published December 12, 1996. Similar methods using *N*-trifluoroacetyl-aminoalkanols as phosphate protecting groups has also been reported by Wilk et al., *J. Org. Chem.*, 62, 6712-6713, 1997. This deprotection is governed by a mechanism that involves removal of *N*-trifluoroacetyl group followed by cyclization of aminoalkyl phosphotriesters to azacyclanes, which is accompanied by the release of the phosphodiester group.

Solid phase techniques continue to play a large role in oligonucleotidic synthetic approaches. Typically, the 3'-most nucleoside is anchored to a solid support which is functionalized with hydroxyl or amino residues. The additional nucleosides are subsequently added in a step-wise fashion to form the desired linkages between the 3'-functional group of the incoming nucleoside, and the 5'-hydroxyl group of the support bound nucleoside. Implicit to this step-wise assembly is the judicious choice of suitable phosphorus protecting groups. Such protecting groups serve to shield phosphorus moieties of the nucleoside base portion of the growing oligomer until such time that it is cleaved from the solid support. Consequently, new protecting groups, which are versatile in oligonucleotidic synthesis, are needed.

A variety of modifications to naturally occurring oligonucleotides have been proposed. Such modifications include labeling with nonisotopic labels, e.g. fluorescein, biotin, digoxigenin, alkaline phosphatase, or other reporter molecules. Other modifications have been made to the ribose phosphate backbone to increase the nuclease stability of the resulting analog. Examples of such modifications include incorporation of methyl phosphonate, phosphorothioate, or phosphorodithioate linkages, and 2'-O-methyl ribose sugar units. Further modifications include those made to modulate uptake and cellular distribution. With the success of these compounds

5 for both diagnostic and therapeutic uses, there exists an ongoing demand for improved oligonucleotides and their analogs.

It is well known that most of the bodily states in multicellular organisms, including most disease states, are effected by proteins. Such proteins, either acting directly or through their enzymatic or other functions, contribute in major proportion to many diseases and regulatory  
10 functions in animals and man. For disease states, classical therapeutics has generally focused upon interactions with such proteins in efforts to moderate their disease-causing or disease-potentiating functions. In newer therapeutic approaches, modulation of the actual production of such proteins is desired. By interfering with the production of proteins, the maximum therapeutic effect may be obtained with minimal side effects. It is therefore a general object of  
15 such therapeutic approaches to interfere with or otherwise modulate gene expression, which would lead to undesired protein formation.

One method for inhibiting specific gene expression is with the use of oligonucleotides, especially oligonucleotides which are complementary to a specific target messenger RNA (mRNA) sequence. Several oligonucleotides are currently undergoing clinical trials for such use.  
20 Phosphorothioate oligonucleotides are presently being used as such antisense agents in human clinical trials for various disease states, including use as antiviral agents. Other mechanisms of action have also been proposed.

Transcription factors interact with double-stranded DNA during regulation of transcription. Oligonucleotides can serve as competitive inhibitors of transcription factors to  
25 modulate their action. Several recent reports describe such interactions (see Bielinska, A., et. al., *Science*, 1990, 250, 997-1000; and Wu, H., et. al., *Gene*, 1990, 89, 203-209).

In addition to such use as both indirect and direct regulators of proteins, oligonucleotides and their analogs also have found use in diagnostic tests. Such diagnostic tests can be performed using biological fluids, tissues, intact cells or isolated cellular components. As with gene  
30 expression inhibition, diagnostic applications utilize the ability of oligonucleotides and their analogs to hybridize with a complementary strand of nucleic acid. Hybridization is the sequence specific hydrogen bonding of oligomeric compounds via Watson-Crick and/or Hoogsteen base

5 pairs to RNA or DNA. The bases of such base pairs are said to be complementary to one another.

Oligonucleotides and their analogs are also widely used as research reagents. They are useful for understanding the function of many other biological molecules as well as in the preparation of other biological molecules. For example, the use of oligonucleotides and their  
 10 analogs as primers in PCR reactions has given rise to an expanding commercial industry. PCR has become a mainstay of commercial and research laboratories, and applications of PCR have multiplied. For example, PCR technology now finds use in the fields of forensics, paleontology, evolutionary studies and genetic counseling. Commercialization has led to the development of kits which assist non-molecular biology-trained personnel in applying PCR. Oligonucleotides  
 15 and their analogs, both natural and synthetic, are employed as primers in such PCR technology.

Oligonucleotides and their analogs are also used in other laboratory procedures. Several of these uses are described in common laboratory manuals such as *Molecular Cloning, A Laboratory Manual*, Second Ed., J. Sambrook, et al., Eds., Cold Spring Harbor Laboratory Press, 1989; and *Current Protocols In Molecular Biology*, F. M. Ausubel, et al., Eds., Current  
 20 Publications, 1993. Such uses include as synthetic oligonucleotide probes, in screening expression libraries with antibodies and oligomeric compounds, DNA sequencing, *in vitro* amplification of DNA by the polymerase chain reaction, and in site-directed mutagenesis of cloned DNA. See Book 2 of *Molecular Cloning, A Laboratory Manual, supra*. See also "DNA-protein interactions and The Polymerase Chain Reaction" in Vol. 2 of *Current Protocols In*  
 25 *Molecular Biology, supra*.

Oligonucleotides and their analogs can be synthesized to have customized properties that can be tailored for desired uses. Thus a number of chemical modifications have been introduced into oligomeric compounds to increase their usefulness in diagnostics, as research reagents and as therapeutic entities. Such modifications include those designed to increase binding to a target  
 30 strand (i.e. increase their melting temperatures,  $T_m$ ), to assist in identification of the oligonucleotide or an oligonucleotide-target complex, to increase cell penetration, to stabilize against nucleases and other enzymes that degrade or interfere with the structure or activity of the

5 oligonucleotides and their analogs, to provide a mode of disruption (terminating event) once sequence-specifically bound to a target, and to improve the pharmacokinetic properties of the oligonucleotide.

For example, antisense oligonucleotides have been modified to be conjugated with lipophilic molecules. The presence of the lipophilic conjugate has been shown to improve  
10 cellular permeation of the oligonucleotide and, accordingly, improve distribution of the oligonucleotide in cells. Further, oligonucleotides conjugated with lipophilic molecules are able to enhance the free uptake of the oligonucleotides without the need for any transfection agents in cell culture studies. Conjugated oligonucleotides are also able to improve the protein binding of oligonucleotides containing phosphodiester linkages.

15 Recently, oligonucleotides selectively labeled with two different reporter groups have attained a widespread interest due to their unique properties. For example, if the reporter groups are paired as a donor and an acceptor of fluorescent energy, a fluorescence resonance energy transfer (FR T) between the two groups may occur. When the distance between the donor and acceptor groups is short, quenching of the fluorescence is observed. In contrast, when the donor and acceptor  
20 are sufficiently far apart, a fluorescent signal is observed. Such a phenomenon has been used to detect formation of a complex between a suitably labeled oligonucleotide and a complementary target nucleic acid. When the oligonucleotide is uncomplexed, the donor and the acceptor groups are sufficiently close so that no fluorescence is detected. However, when the oligonucleotide complexes with the nucleic acid, the donor and acceptor groups are forced to move away from each  
25 other, thereby restoring the fluorescence signal. Oligonucleotides that exhibit such a hybridization-dependent fluorescence have been termed "molecular beacons." Molecular beacons may be useful in numerous applications, such as real-time monitoring of hybridization in PCR, in molecular biosensors, on surfaces, in blood, in living cells, and *in vivo*. Synthesis of double labeled oligonucleotides has been performed with the aid of dye-labeled solid supports and  
30 phosphoramidites. In addition, oligonucleotides that bear an amino and an activated thiol group at opposite ends have been synthesized using modified phosphoramidites and solid supports, where chemoselective labeling is performed post-synthetically. However, the known methods for

5 synthesizing double labeled oligonucleotides are restricted to the use of only certain dyes that are available as phosphoramidite, solid support, and chemoselective reagents.


In light of the foregoing, there is a continued need for selectively labeled oligonucleotides, methods for making the oligonucleotides, and compounds useful therefor. The labeled oligonucleotides should provide improved cellular permeation, enhanced free uptake of the  
10 oligonucleotide in cell culture studies, and improved protein binding, especially for oligonucleotides containing phosphodiester linkages. In addition, there is a need for methods for producing oligonucleotides selectively labeled at one or more of the 3'-terminal nucleotide, 5'-terminal nucleotide, and internucleotides with one or more different conjugate groups. The methods should also provide for such labeled oligonucleotides without the need for post-synthetic labeling.

15

## SUMMARY OF THE INVENTION

The present invention allows for the selective functionalization of oligonucleotides with conjugate groups. In particular, the oligonucleotides can be selectively functionalized with a first conjugate group at the 3'-terminal nucleotide and optionally functionalized with a second conjugate  
20 group at the 5'-terminal nucleotide and/or one or more internucleotides. Alternatively, the oligonucleotides can be selectively functionalized with a first conjugate group at the 5'-terminal nucleotide and optionally functionalized with a second conjugate group at one or more internucleotides. In yet another embodiment, the oligonucleotides can be functionalized with a first conjugate group at one or more internucleotides and with a second conjugate group at one or more  
25 different internucleotides.


It is an object of the present invention to provide oligonucleotides having the formula:



5

wherein:

$R_1$  is hydroxyl, a protected hydroxyl or a group having the formula:




10

$\text{Q}_0$  is O or S;

$\text{R}_4$  is  $\text{O}^-$ , hydroxyl or a protected hydroxyl;

$\text{R}_2$  is hydroxyl, a protected hydroxyl or a group having the formula:



15

each  $\text{R}_3$  is H, a 2'-substituent group or a protected 2'-substituent group;

each X is, independently,  $\text{O}^-$ , hydroxyl, protected hydroxyl or  $-\text{S}-\text{L}_3$ ;

each Bx is an optionally protected heterocyclic base moiety;

n is from 3 to about 50; and

5 L<sub>1</sub>, L<sub>2</sub> and each of said L<sub>3</sub> are, independently, a conjugate group

It is a further object of the present invention to provide methods for the preparation of selectively functionalized oligonucleotides having the above formula.

10 It is yet a further object of the present invention to provide synthetic intermediates useful in such methods.

Other objects will be apparent to those skilled in the art.

### BRIEF DESCRIPTION OF THE DRAWINGS

15 The advantageous features of the invention will be more fully appreciated when considered based on the following detailed description and the attached drawings, wherein:

FIG. 1 is an RP HPLC profile of compound 120 in its crude form prior to removal of the DMT group;

FIG. 2 is an RP HPLC profile of compound 120 in its crude form after removal of the DMT group;

20 FIG. 3 is an RP HPLC profile of compound 120 in its final form;

FIG. 4 is an RP HPLC profile of compound 129 in its crude form prior to removal of the DMT group;

FIG. 5 is an RP HPLC profile of compound 129 in its crude form after removal of the DMT group;

25 FIG. 6 is an RP HPLC profile of compound 129 in its final form;

FIG. 7 is an RP HPLC profile of compound 126 in its crude form after removal of the DMT group;

FIG. 8 is an RP HPLC profile of compound 126 in its final form;

30 FIG. 9 is an RP HPLC profile of compound 121 in its crude form after removal of the DMT group; and

FIG. 10 is an RP HPLC profile of compound 121 in its final form.



## 5 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS


The present invention provides methods for preparing compounds that comprise a plurality of linked nucleotides wherein the linked nucleotides are selectively functionalized or labeled with conjugate groups. The compounds can be functionalized at the 3'-terminal nucleotide, at the 5'-terminal nucleotide, at an internucleotide, at the 3'-terminal nucleotide with a first conjugate group and at the 5'-terminal nucleotide with a second conjugate group, at the 3'-terminal nucleotide with a first conjugate group and at one or more internucleotides with a second conjugate group, at the 5'-terminal nucleotide with a first conjugate group and at one or more internucleotides with a second conjugate group, or at one or more internucleotides with a first conjugate group and one or more internucleotides with a second conjugate group.

As used herein, "oligomer" and "oligomeric compound" refer to compounds containing a plurality of monomeric subunits that are joined by phosphorus-containing linkages, such as phosphite, phosphodiester, phosphorothioate, and/or phosphorodithioate linkages. Oligomeric compounds therefore include oligonucleotides, their analogs, and synthetic oligonucleotides. The methods of the invention are used for the preparation of oligonucleotides and their analogs.

As used herein, the term "oligonucleotide analog" means compounds that can contain both naturally occurring (i.e. "natural") and non-naturally occurring synthetic moieties, for example, nucleosidic subunits containing modified sugar and/or nucleobase portions. Such oligonucleotide analogs are typically structurally distinguishable from, yet functionally interchangeable with, naturally occurring or synthetic wild type oligonucleotides. Thus, oligonucleotide analogs include all such structures which function effectively to mimic the structure and/or function of a desired RNA or DNA strand, for example, by hybridizing to a target. The term synthetic nucleoside, for the purpose of the present invention, refers to a modified nucleoside. Representative modifications include modification of a heterocyclic base portion of a nucleoside to give a non-naturally occurring nucleobase, a sugar portion of a nucleoside, or both simultaneously.


The present invention relates to processes for preparing an oligonucleotide having the formula:

5



wherein:


$R_1$  is hydroxyl, a protected hydroxyl or a group having the formula:



$Q_0$  is O or S;

$R_4$  is O<sup>-</sup>, hydroxyl or protected hydroxyl;

$R_2$  is hydroxyl, a protected hydroxyl or a group having the formula:



each  $R_3$  is H, a 2'-substituent group or a protected 2'-substituent group;

each X is, independently, O or -S- $L_3$ ;

each Bx is an optionally protected heterocyclic base moiety;

5           n is from 3 to about 50; and  
           $L_1$ ,  $L_2$  and each of said  $L_3$  are, independently, a conjugate group.

          In one embodiment,  $R_1$  comprises a first conjugate group and  $R_2$  optionally comprises a second conjugate group. In another embodiment,  $R_1$  comprises a first conjugate group and one  
10 or more  $R_3$  optionally comprise a second conjugate group. In yet another embodiment,  $R_2$  comprises a first conjugate group and one or more  $R_3$  optionally comprise a second conjugate group. In still another embodiment, one or more  $R_3$  comprise a first conjugate group and one or more  $R_3$  optionally comprise a second conjugate group.

          Oligonucleotides, or more broadly oligomeric compounds, according to the present invention  
15 preferably comprise from about 3 to about 50 nucleosides. It is more preferred that such compounds comprise from about 8 to about 30 nucleosides, with 15 to 25 nucleosides being particularly preferred. When used as "building blocks" in assembling larger oligomeric compounds (i.e., as synthons), smaller oligomeric compounds are preferred. Libraries of dimeric, trimeric, or higher order compounds can be prepared for use as synthons in the methods of the invention. The use of  
20 small sequences synthesized via solution phase chemistries in automated synthesis of larger oligonucleotides enhances the coupling efficiency and the purity of the final oligonucleotides. See for example: Miura, K., *et al.*, *Chem. Pharm. Bull.*, **1987**, 35, 833-836; Kumar, G., and Poonian, M.S., *J. Org. Chem.*, **1984**, 49, 4905-4912; Bannwarth, W., *Helvetica Chimica Acta*, **1985**, 68, 1907-1913; Wolter, A., *et al.*, *nucleosides and nucleotides*, **1986**, 5, 65-77, each of which are hereby  
25 incorporated by reference in their entirety.

          The attachment of conjugate groups to oligonucleotides and analogs thereof is well documented in the prior art. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, poly-  
30 ethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, acridine,

5 fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion.

10 Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992, United States Patent No. 5,578,718, issued July 1, 1997, and United States Patent No. 5,218,105. Each of the foregoing is commonly assigned with this application. The entire disclosure of each is incorporated herein by reference.

Preferred conjugate groups amenable to the present invention include lipid moieties such  
15 as a cholesterol moiety (Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553), cholic acid (Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053), a thioether, *e.g.*, hexyl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992, 660, 306; Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765), a thiocholesterol (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20, 533), an aliphatic chain, *e.g.*, dodecandiol or undecyl residues (Saison-Behmoaras *et al.*, *EMBO J.*, 1991,  
20 10, 111; Kabanov *et al.*, *FEBS Lett.*, 1990, 259, 327; Svinarchuk *et al.*, *Biochimie*, 1993, 75, 49), a phospholipid, *e.g.*, di-hexadecyl-*rac*-glycerol or triethylammonium-1,2-di-O-hexadecyl-*rac*-glycero-3-H-phosphonate (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36, 3651; Shea *et al.*, *Nucl. Acids Res.*, 1990, 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, *Nucleosides & Nucleotides*, 1995, 14, 969), adamantane acetic acid (Manoharan *et al.*,  
25 *Tetrahedron Lett.*, 1995, 36, 3651), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264, 229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, 277, 923).

Other groups for modifying antisense properties include RNA cleaving complexes, pyrenes, metal chelators, porphyrins, alkylators, hybrid intercalator/ligands and photo-  
30 crosslinking agents. RNA cleavers include *o*-phenanthroline/Cu complexes and Ru(bipyridine)<sub>3</sub><sup>2+</sup> complexes. The Ru(bpy)<sub>3</sub><sup>2+</sup> complexes interact with nucleic acids and cleave nucleic acids photochemically. Metal chelators are include EDTA, DTPA, and *o*-phenanthroline.

5 Alkylators include compounds such as iodoacetamide. Porphyrins include porphine, its substituted forms, and metal complexes. Pyrenes include pyrene and other pyrene-based carboxylic acids that could be conjugated using the similar protocols.


Hybrid intercalator/ligands include the photonuclease/intercalator ligand 6-[[[9-[[6-(4-nitrobenzamido)hexyl]amino]acridin-4-yl]carbonyl]amino]hexanoyl-pentafluorophenyl ester.

10 This compound has two noteworthy features: an acridine moiety that is an intercalator and a p-nitro benzamido group that is a photonuclease.

Photo-crosslinking agents include aryl azides such as, for example, N-hydroxysuccinimidyl-4-azidobenzoate (HSAB) and N-succinimidyl-6(-4'-azido-2'-nitrophenyl-amino)hexanoate (SANPAH). Aryl azides conjugated to oligonucleotides effect crosslinking  
15 with nucleic acids and proteins upon irradiation. They also crosslink with carrier proteins (such as KLH or BSA), raising antibody against the oligonucleotides.

Vitamins according to the invention generally can be classified as water soluble or lipid soluble. Water soluble vitamins include thiamine, riboflavin, nicotinic acid or niacin, the vitamin B<sub>6</sub> pyridoxal group, pantothenic acid, biotin, folic acid, the B<sub>12</sub> cobamide coenzymes, inositol,  
20 choline and ascorbic acid. Lipid soluble vitamins include the vitamin A family, vitamin D, the vitamin E tocopherol family and vitamin K (and phytols). The vitamin A family, including retinoic acid and retinol, are absorbed and transported to target tissues through their interaction with specific proteins such as cytosol retinol-binding protein type II (CRBP-II), retinol-binding protein (RBP), and cellular retinol-binding protein (CRBP). These proteins, which have been found in various parts  
25 of the human body, have molecular weights of approximately 15 kD. They have specific interactions with compounds of vitamin-A family, especially, retinoic acid and retinol.

The conjugate groups, L<sub>1</sub>, L<sub>2</sub>, and/or L<sub>3</sub>, are optionally attached to the oligonucleotides of the present invention through a linking group. Suitable linking groups include, but are not limited to, dialkylglycerol linkers. Preferred dialkylglycerol linkers have the structure:



where L is L<sub>1</sub>, L<sub>2</sub>, or L<sub>3</sub>.


As used herein, the term "2'-substituent group" refers to groups that are attached to selected sugar moieties at the 2'-position. However, substituent groups can alternatively be attached to other positions of the sugar moieties (e.g., the 3'- and/or 5'-positions), selected heterocyclic base moieties, or at both the heterocyclic base and the sugar moiety.

A representative list of substituent groups amenable to the present invention include hydrogen, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>5</sub>-C<sub>20</sub> aryl, O-alkyl, O-alkenyl, O-alkynyl, O-alkylamino, O-alkylalkoxy, O-alkylaminoalkyl, O-alkyl imidazole, S-alkyl, S-alkenyl, S-alkynyl, NH-alkyl, NH-alkenyl, NH-alkynyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, N-phthalimido, halogen (particularly fluoro), amino, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalators, reporter groups, conjugates, polyamine, polyamide, polyalkylene glycol, and polyethers of the formula (O-alkyl)<sub>m</sub>, where m is 1 to about 10. Preferred among these polyethers are linear and cyclic polyethylene glycols (PEGs), and (PEG)-containing groups, such as crown ethers and those which are disclosed by Ouchi *et al.* (*Drug Design and Discovery* **1992**, 9, 93), Ravasio *et al.* (*J. Org. Chem.* **1991**, 56, 4329) and Delgado *et al.* (*Critical Reviews in Therapeutic Drug Carrier Systems* **1992**, 9, 249), each of which is herein incorporated by reference in its entirety. Further sugar modifications are disclosed in Cook, P.D., *Anti-Cancer Drug Design*, **1991**, 6, 585-607. Fluoro, O-alkyl, O-alkylamino, O-alkyl imidazole, O-alkylaminoalkyl, and alkyl amino substitution is described in United States Patent Application serial number 08/398,901, filed March 6, 1995, entitled Oligomeric Compounds having Pyrimidine Nucleotide(s) with 2' and 5' Substitutions, hereby incorporated by reference in

5 its entirety.

Additional substituent groups amenable to the present invention include -SR and -NR<sub>2</sub> groups, wherein each R is, independently, hydrogen, a protecting group or substituted or unsubstituted alkyl, alkenyl, or alkynyl. 2'-SR nucleosides are disclosed in United States Patent No. 5,670,633, issued September 23, 1997, hereby incorporated by reference in its entirety. The  
 10 incorporation of 2'-SR monomer synthons are disclosed by Hamm *et al.*, *J. Org. Chem.*, **1997**, 62, 3415-3420. 2'-NR<sub>2</sub> nucleosides are disclosed by Goettingen, M., *J. Org. Chem.*, **1996**, 61, 6273-6281; and Polushin *et al.*, *Tetrahedron Lett.*, **1996**, 37, 3227-3230.

Further representative substituent groups can include groups having the structure of one of formula I or II:




wherein:

Z<sub>0</sub> is O, S or NH;

J is a single bond, O or C(=O);

E is C<sub>1</sub>-C<sub>10</sub> alkyl, N(R<sub>5</sub>)(R<sub>6</sub>), N(R<sub>5</sub>)(R<sub>7</sub>), N=C(R<sub>5</sub>)(R<sub>6</sub>), N=C(R<sub>5</sub>)(R<sub>7</sub>) or has one of formula  
 20 III or IV;



5

each  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$  and  $R_{12}$  is, independently, hydrogen,  $C(O)R_{13}$ , substituted or unsubstituted  $C_1$ - $C_{10}$  alkyl, substituted or unsubstituted  $C_2$ - $C_{10}$  alkenyl, substituted or unsubstituted  $C_2$ - $C_{10}$  alkynyl, alkylsulfonyl, arylsulfonyl, a chemical functional group or a conjugate group, wherein the substituent groups are selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl;

or optionally,  $R_9$  and  $R_{10}$ , together form a phthalimido moiety with the nitrogen atom to which they are attached;

or optionally,  $R_{11}$  and  $R_{12}$ , together form a phthalimido moiety with the nitrogen atom to which they are attached;

each  $R_{13}$  is, independently, substituted or unsubstituted  $C_1$ - $C_{10}$  alkyl, trifluoromethyl, cyanoethoxy, methoxy, ethoxy, t-butoxy, allyloxy, 9-fluorenylmethoxy, 2-(trimethylsilyl)-ethoxy, 2,2,2-trichloroethoxy, benzyloxy, butyryl, iso-butyryl, phenyl or aryl;

$R_5$  is T-L,

T is a bond or a linking moiety;

L is a chemical functional group, a conjugate group or a solid support material;

each  $R_5$  and  $R_6$  is, independently, H, a nitrogen protecting group, substituted or unsubstituted  $C_1$ - $C_{10}$  alkyl, substituted or unsubstituted  $C_2$ - $C_{10}$  alkenyl, substituted or unsubstituted  $C_2$ - $C_{10}$  alkynyl, wherein said substitution is  $OR_3$ ,  $SR_3$ ,  $NH_3^+$ ,  $N(R_{14})(R_{15})$ , guanidino or acyl where said acyl is an acid amide or an ester;

or  $R_5$  and  $R_6$ , together, are a nitrogen protecting group or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

or  $R_5$ , T and L, together, are a chemical functional group;

each  $R_{14}$  and  $R_{15}$  is, independently, H,  $C_1$ - $C_{10}$  alkyl, a nitrogen protecting group, or  $R_{14}$  and  $R_{15}$ , together, are a nitrogen protecting group;

or  $R_{14}$  and  $R_{15}$  are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;



- 5             $Z_4$  is OX, SX, or  $N(X)_2$ ;  
             each X is, independently, H,  $C_1-C_8$  alkyl,  $C_1-C_8$  haloalkyl,  $C(=NH)N(H)R_{16}$ ,  
 $C(=O)N(H)R_{16}$  or  $OC(=O)N(H)R_{16}$ ;  
              $R_{16}$  is H or  $C_1-C_8$  alkyl;  
              $Z_1$ ,  $Z_2$  and  $Z_3$  comprise a ring system having from about 4 to about 7 carbon atoms or  
10    having from about 3 to about 6 carbon atoms and 1 or 2 heteroatoms wherein said heteroatoms  
         are selected from oxygen, nitrogen and sulfur and wherein said ring system is aliphatic,  
         unsaturated aliphatic, aromatic, or saturated or unsaturated heterocyclic;

- $Z_5$  is alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10  
         carbon atoms, alkynyl having 2 to about 10 carbon atoms, aryl having 6 to about 14 carbon  
15    atoms,  $N(R_5)(R_6)OR_5$ , halo,  $SR_5$  or CN;  
             each  $q_1$  is, independently, an integer from 1 to 10;  
             each  $q_2$  is, independently, 0 or 1;  
              $q_3$  is 0 or an integer from 1 to 10;  
              $q_4$  is an integer from 1 to 10;  
20     $q_5$  is from 0, 1 or 2; and  
             provided that when  $q_3$  is 0,  $q_4$  is greater than 1.

Representative substituent groups of Formula I are disclosed in United States Patent  
Application Serial No. 09/130,973, filed August 7, 1998, entitled "Capped 2'-Oxyethoxy  
Oligonucleotides," hereby incorporated by reference in its entirety.

- 25            Representative cyclic substituent groups of Formula II are disclosed in United States  
Patent Application Serial No. 09/123,108, filed July 27, 1998, entitled "RNA Targeted  
2'-Modified Oligonucleotides that are Conformationally Preorganized," hereby incorporated by  
reference in its entirety.

- Particularly preferred sugar substituent groups include  $O[(CH_2)_nO]_mCH_3$ ,  $O(CH_2)_nOCH_3$ ,  
30     $O(CH_2)_nNH_2$ ,  $O(CH_2)_nCH_3$ ,  $O(CH_2)_nONH_2$ , and  $O(CH_2)_nON[(CH_2)_nCH_3]_2$ , where n and m are from  
1 to about 10.

Some preferred oligomeric compounds of the invention contain, at least one nucleoside

5 having one of the following substituent groups: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, poly-alkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligomeric compound, or a group for  
10 improving the pharmacodynamic properties of an oligomeric compound, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy [2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE] (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78, 486), *i.e.*, an alkoxyalkoxy group. A further preferred modification is 2'-dimethylaminooxyethoxy, *i.e.*, a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE.

15 Representative aminooxy substituent groups are described in co-owned United States Patent Application serial number 09/344,260, filed June 25, 1999, entitled "Aminooxy-Functionalized Oligomers"; and United States Patent Application serial number 09/370,541, filed August 9, 1999, entitled "Aminooxy-Functionalized Oligomers and Methods for Making Same;" hereby incorporated by reference in their entirety.

20 Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on nucleosides and oligomers, particularly the 3' position of the sugar on the 3' terminal nucleoside or at a 3'-position of a nucleoside that has a linkage from the 2'-position such as a 2'-5' linked oligomer and at the 5' position of a 5' terminal nucleoside. Oligomers may also have  
25 sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugars structures include, but are not limited to, U.S. Patents 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of  
30 which are commonly owned, and each of which is herein incorporated by reference, and commonly owned United States patent application 08/468,037, filed on June 5, 1995, also herein incorporated by reference.

5 Representative guanidino substituent groups that are shown in formula III and IV are disclosed in co-owned United States Patent Application 09/349,040, entitled "Functionalized Oligomers", filed July 7, 1999, hereby incorporated by reference in its entirety.

Representative acetamido substituent groups are disclosed in United States Patent Application 09/378,568, entitled "2'-O-Acetamido Modified Monomers and Oligomers", filed 10 August 19, 1999, hereby incorporated by reference in its entirety.

Representative dimethylaminoethoxyethyl substituent groups are disclosed in International Patent Application PCT/US99/17895, entitled "2'-O-Dimethylaminoethoxyethyl-Modified Oligonucleotides", filed August 6, 1999, hereby incorporated by reference in its entirety.

15 "B" and "Bx," as used herein, is intended to indicate a heterocyclic base moiety. A heterocyclic base moiety (often referred to in the art simply as a "base" or a "nucleobase") amenable to the present invention includes both naturally and non-naturally occurring nucleobases. The heterocyclic base moiety further may be protected wherein one or more functionalities of the base bears a protecting group. As used herein, "unmodified" or "natural" 20 nucleobases include the purine bases adenine and guanine, and the pyrimidine bases thymine, cytosine and uracil. Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5- 25 uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. 30 Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, *Angewandte*

- 5 *Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993.

Certain heterocyclic base moieties are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention to complementary targets. These include  
10 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2 °C (*Id.*, pages 276-278) and are presently preferred base substitutions, even more particularly when combined with selected 2'-sugar modifications such as 2'-methoxyethyl groups.

15 Representative United States patents that teach the preparation of heterocyclic base moieties (modified nucleobases) include, but are not limited to, U.S. Patents 3,687,808; 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned, and each of which is herein incorporated by  
20 reference, and commonly owned United States patent application 08/762,587, filed on December 10, 1996, also herein incorporated by reference.


The present invention provides oligomeric compounds comprising a plurality of linked nucleosides wherein the preferred internucleoside linkage is a 3',5'-linkage. Alternatively, however, 2',5'-linkages can be used (as described in U.S. Application Serial No. 09/115,043,  
25 filed July 14, 1998). A 2',5'-linkage is one that covalently connects the 2'-position of the sugar portion of one nucleotide subunit with the 5'-position of the sugar portion of an adjacent nucleotide subunit.

The compounds described herein may have asymmetric centers. Unless otherwise indicated, all chiral, diastereomeric, and racemic forms are included in the present invention.  
30 Geometric isomers may also be present in the compounds described herein, and all such stable isomers are contemplated by the present invention. It will be appreciated that compounds in accordance with the present invention that contain asymmetrically substituted carbon atoms may

- 5 be isolated in optically active or racemic forms or by synthesis.


The present invention includes all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of example, and without limitation, isotopes of hydrogen include tritium and deuterium.

- 10 In one embodiment, the present invention provides a process for preparing an oligonucleotide having the formula:



- 15 wherein:


$R_1$  is hydroxyl, a protected hydroxyl or a group having the formula:



$Q_0$  is O or S;

$R_4$  is O<sup>-</sup>, hydroxyl, or a protected hydroxyl;

- 20  $R_2$  is hydroxyl, a protected hydroxyl or a group having the formula:



each  $R_3$  is H, a 2'-substituent group or a protected 2'-substituent group;

each X is, independently, O<sup>-</sup>, hydroxyl, protected hydroxyl, or -S-L<sub>3</sub>;

each B<sub>x</sub> is an optionally protected heterocyclic base moiety;


n is from 3 to about 50; and

L<sub>1</sub>, L<sub>2</sub> and each of said L<sub>3</sub> are, independently, a ligand.


In preferred embodiments, the oligonucleotides have at least two different ligands attached covalently thereto. Further, it is particularly preferred that L<sub>1</sub> is different than L<sub>2</sub>, L<sub>2</sub> is different than each of L<sub>1</sub> and L<sub>3</sub>, and each L<sub>3</sub> attached at a particular position is different from a second L<sub>3</sub> attached at a different position. In addition, said two different ligands covalently attached thereto are preferably positioned to said oligonucleotide at L<sub>1</sub> and L<sub>2</sub>, L<sub>1</sub> and L<sub>3</sub>, L<sub>2</sub> and L<sub>3</sub>, or two of said L<sub>3</sub> groups.

The process comprises the steps of:

a) providing a derivatized solid support for oligonucleotide synthesis, said derivatized solid support being derivatized with a group having one of the structures:



or




wherein

T is a bifunctional linking moiety linked to the solid support; and

- 5             $Q_1$  is an acid labile hydroxyl protecting group;
- b)            treating said solid support with an acidic reagent to deblock said acid labile hydroxyl protecting group to give a free hydroxyl group;
- c)            reacting said free hydroxyl group with a phosphoramidite composition to form an extended compound, said phosphoramidite composition having the formula:


10



wherein

- 15             $Q_2$  is a 5'-terminal acid labile hydroxyl protecting group;
- $Q_3$  is a phosphorus protecting group; and
- $Z_6$  and  $Z_7$  are, independently,  $C_{1-6}$  alkyl;
- or  $Z_6$  and  $Z_7$  are joined together to form a 4- to 7-membered heterocyclic ring system including the nitrogen atom to which  $Z_6$  and  $Z_7$  are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S;
- 20            d)            optionally treating said extended compound with a capping agent to form a capped compound;
- e)            optionally oxidizing said capped compound to form an oxidized compound;
- f)            repeating steps b) through e) at least three times to form a further extended
- 25    compound;
- g)            optionally treating said further extended compound with an acidic reagent effective to deblock said acid labile hydroxyl protecting group to give a free hydroxyl group and

- 5 reacting said free hydroxyl group of said further extended compound with a compound of the formula:



- 10 thereby forming a 5'-functionalized further extended compound, wherein Q<sub>5</sub> is an acid labile hydroxyl protecting group;

h) treating said further extended compound or said 5'-functionalized further extended compound for a time and under conditions effective to remove at least one phosphorus protecting group giving at least one deblocked phosphorothioate linkage;

- 15 i) reacting said deblocked phosphorothioate linkage with a ligand that is reactive with and forms a covalent bond with said deblocked phosphorothioate linkage; and

j) optionally repeating steps h) and i) to give said oligonucleotide.

- 20 The methods of the present invention are useful for the preparation of all compounds containing phosphorus functionalities. As used herein, functionality includes, but is not limited to phosphite, phosphodiester, phosphorothioate, and/or phosphorodithioate residues, and oligomeric compounds containing monomeric subunits that are joined by a variety of functionality linkages, including phosphite, phosphodiester, phosphorothioate, and/or phosphorodithioate linkages.

- 25 The oligomeric compounds in accordance with the invention can be used in diagnostics, therapeutics and as research reagents and kits. They can be used in pharmaceutical compositions by including a suitable pharmaceutically acceptable diluent or carrier. They further can be used



5 for treating organisms having a disease characterized by the undesired production of a protein. The organism should be contacted with an oligonucleotide having a sequence that is capable of specifically hybridizing with a strand of nucleic acid coding for the undesirable protein. Treatments of this type can be practiced on a variety of organisms ranging from unicellular prokaryotic and eukaryotic organisms to multicellular eukaryotic organisms. Any organism that  
10 utilizes DNA-RNA transcription or RNA-protein translation as a fundamental part of its hereditary, metabolic or cellular control is susceptible to therapeutic and/or prophylactic treatment in accordance with the invention. Seemingly diverse organisms such as bacteria, yeast, protozoa, algae, all plants and all higher animal forms, including warm-blooded animals, can be treated. Further, each cell of multicellular eukaryotes can be treated, as they include both DNA-  
15 RNA transcription and RNA-protein translation as integral parts of their cellular activity. Furthermore, many of the organelles (*e.g.*, mitochondria and chloroplasts) of eukaryotic cells also include transcription and translation mechanisms. Thus, single cells, cellular populations or organelles can also be included within the definition of organisms that can be treated with therapeutic or diagnostic oligonucleotides.

20 The reactions of the synthetic methods claimed herein are carried out in suitable solvents which may be readily understood by those skilled in the art of organic synthesis, the suitable solvents generally being any solvent which is substantially nonreactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, *i.e.*, temperatures may range from the solvent's freezing temperature to the solvent's  
25 boiling temperature. A given reaction may be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step may be selected.

Methods for assembling oligomers in accordance with the present invention include both solution phase and solid phase chemistries. Representative solution phase techniques are  
30 described in United States Patent No. 5,210,264, which is assigned to the assignee of the present invention. In preferred embodiments, the methods of the present invention are employed for use in iterative solid phase oligonucleotide synthetic regimes. Representative solid phase techniques

are those typically employed for DNA and RNA synthesis utilizing standard phosphoramidite chemistry, (*see, e.g.*, Protocols For Oligonucleotides And Analogs, Agrawal, S., ed., Humana Press, Totowa, NJ, 1993, hereby incorporated by reference in its entirety).

Solid supports according to the invention include those generally known in the art to be suitable for use in solid phase methodologies, including, for example, controlled pore glass (CPG), oxalyl-controlled pore glass (*see, e.g.*, Alul, *et al.*, *Nucleic Acids Research* **1991**, *19*, 1527, hereby incorporated by reference in its entirety), TentaGel Support -- an aminopolyethyleneglycol derivatized support (*see, e.g.*, Wright, *et al.*, *Tetrahedron Letters* **1993**, *34*, 3373, hereby incorporated by reference in its entirety) and Poros -- a copolymer of polystyrene/ divinylbenzene.

In a preferred embodiment, the solid support is derivatized to provide an acid labile trialkoxytrityl group, such as a trimethoxytrityl group (TMT). Without being bound by theory, it is expected that the presence of the trialkoxytrityl protecting group will permit initial detritylation under conditions commonly used on DNA synthesizers. For a faster release of oligonucleotide material in solution with aqueous ammonia, a diglycoate linker is optionally introduced onto the support.


A preferred synthetic solid phase synthesis utilizes phosphoramidites as activated phosphate compounds. In this technique, a phosphoramidite monomer is reacted with a free hydroxyl on the growing oligomer chain to produce an intermediate phosphite compound, which is subsequently oxidized to the P<sup>V</sup> state using standard methods. This technique is commonly used for the synthesis of several types of linkages including phosphodiester, phosphorothioate, and phosphorodithioate linkages.

Typically, the first step in such a process is attachment of a first monomer or higher order subunit to a solid support using standard methods and procedures known in the art. Solid supports are substrates which are capable of serving as the solid phase in solid phase synthetic methodologies, such as those described in Caruthers U.S. Patents Nos. 4,415,732; 4,458,066; 4,500,707; 4,668,777; 4,973,679; and 5,132,418; and Koster U.S. Patents Nos. 4,725,677 and Re. 34,069. A linker is optionally positioned between the terminal nucleotide and the solid support.

- 5 Linkers are known in the art as short molecules which serve to connect a solid support to functional groups (e.g., hydroxyl groups) of initial synthon molecules in solid phase synthetic techniques. Suitable linkers are disclosed in, for example, *Oligonucleotides And Analogues A Practical Approach*, Ekstein, F. Ed., IRL Press, N.Y, 1991, Chapter 1, pages 1-23, hereby incorporated by reference in its entirety.
- 10 The support-bound monomer or higher order synthon is then treated to remove the protecting group from the free terminal end. Typically, this is accomplished by treatment with acid. The solid support bound monomer, or higher order oligomer, is then reacted with individual monomeric or higher order building blocks (i.e., synthons) to form a compound which has a phosphite or thiophosphite linkage. In preferred embodiments, the synthons reacted under
- 15 anhydrous conditions in the presence of an activating agent such as, for example, 1H-tetrazole, 5-(4-nitrophenyl)-1H-tetrazole, or diisopropylamino tetrazolide.

In one preferred embodiment, the oligonucleotides are assembled according to the method described in U.S. Patent No. 6,121,437, which is hereby incorporated by reference in its entirety. Accordingly, the phosphate groups of internucleosidic nucleotides that are to be

20 functionalized are protected with derivatives of 2-benzamidoethyl groups. In particular, the oligonucleotides are formed by reacting a compound of Formula V:



V

5

wherein:

W is selected independently from O and S;

X is selected independently from O and S;

Y is selected independently from O and NR<sup>22</sup>;

10

Z is selected independently from a single bond, O, and NR<sup>22</sup>;

R<sup>21</sup>, at each occurrence, is selected independently from C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, CN, NO<sub>2</sub>, Cl, Br, F, I, CF<sub>3</sub>, OR<sup>23</sup>, NR<sup>24</sup>R<sup>25</sup>, and phenyl;

alternatively, two R<sup>21</sup> groups, when on adjacent carbons of the phenyl ring, join to form a naphthyl ring that includes said phenyl ring;

15

R<sup>22</sup>, at each occurrence, is selected independently from H, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, and phenyl;

R<sup>20</sup>, at each occurrence, is selected independently from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, and phenyl;

R<sup>23</sup> is selected independently from C<sub>1-6</sub> alkyl, C<sub>3-6</sub> cycloalkyl, and phenyl;

20

R<sup>24</sup> and R<sup>25</sup>, at each occurrence, are selected independently from C<sub>1-6</sub> alkyl, C<sub>3-6</sub> cycloalkyl, and phenyl;

n is selected independently from 0, 1, 2, and 3; and


m is selected independently from 0, 1, 2, and 3;

25


R<sup>17</sup>, at each occurrence, is selected independently from H, a hydroxyl protecting group, and a linker connected to a solid support;

R<sup>18</sup>, at each occurrence, is independently H, hydroxyl, C<sub>1-20</sub> alkyl, C<sub>3-20</sub> alkenyl, C<sub>2-20</sub> alkynyl, halogen, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, N-phthalimido, imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, or one of formula VI or VII:

30



VI



VII

wherein

E is selected from  $\text{C}_{1-10}$  alkyl,  $\text{N}(\text{R}^{26})(\text{R}^{27})$  and  $\text{N}=\text{C}(\text{R}^{26})(\text{R}^{27})$ ;

$\text{R}^{26}$  and  $\text{R}^{27}$  are independently selected from H,  $\text{C}_{1-10}$  alkyl, dialkylaminoalkyl, a nitrogen protecting group, a tethered or untethered conjugate group, a linker to a solid support, or alternatively  $\text{R}^{26}$  and  $\text{R}^{27}$ , together, are joined in a nitrogen protecting group or a ring structure that can include at least one additional heteroatom selected from N and O;

$q^1$  is from 1 to 10;

$q^2$  is from 1 to 10;

$q^3$  is 0 or 1;

$\text{Z}_{11}$  is  $\text{OR}^{28}$ ,  $\text{SR}^{28}$ , or  $\text{N}(\text{R}^{28})_2$ ;

$\text{R}^{28}$  is selected independently from H,  $\text{C}_1\text{--C}_8$  alkyl,  $\text{C}_1\text{--C}_8$  haloalkyl,  $\text{C}(=\text{NH})\text{N}(\text{H})\text{R}^{29}$ ,  $\text{C}(=\text{O})\text{N}(\text{H})\text{R}^{29}$  and  $\text{OC}(=\text{O})\text{N}(\text{H})\text{R}^{29}$ ;

$\text{R}^{29}$  is H or  $\text{C}_1\text{--C}_8$  alkyl;

$\text{Z}_8$ ,  $\text{Z}_9$  and  $\text{Z}_{10}$  comprise a ring system having from about 4 to about 7 carbon atoms or having from about 3 to about 6 carbon atoms and 1 or 2 heteroatoms wherein said heteroatoms are selected from oxygen, nitrogen and sulfur and wherein said ring system is aliphatic, unsaturated aliphatic, aromatic, or saturated or unsaturated heterocyclic;


$\text{Z}_{11}$  is alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, aryl having 6 to about 14 carbon atoms,  $\text{N}(\text{R}^{26})(\text{R}^{27})$ ,  $\text{OR}^{26}$ , halo,  $\text{SR}^{26}$  or CN; and

$q^4$  is, 0, 1 or 2;


$\text{R}^{19}$  is selected independently from  $\text{NR}^{30}\text{R}^{31}$ , and a 5-6 membered heterocyclic system containing 1-4 heteroatoms selected independently from N, O, and S;

$\text{R}^{30}$  and  $\text{R}^{31}$ , at each occurrence, are selected independently from  $\text{C}_{1-10}$  alkyl,  $\text{C}_{3-7}$  cycloalkyl, and isopropyl;

- 5  $X^1$  is selected independently from O and S;  
 B, at each occurrence, is independently selected from a protected or unprotected naturally occurring nucleobase, and a protected or unprotected non-naturally occurring nucleobase;  
 q is selected independently from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;  
 10 p is an integer selected independently from 0 to about 50;  
 Q, at each occurrence, is selected independently from OH, SH, and



- 15 with a compound of the following formula:




wherein:

- $R^{32}$  is selected independently from a hydroxyl protecting group, and a linker  
 20 connected to a solid support; and  
 p' is an integer selected independently from 0 to about 50.

In another embodiment, the oligonucleotides are assembled from building blocks having the formula VIII:

5



## VIII

wherein:

DMTr is a 4,4'- dimethoxytrityl group;

iPr is an isopropyl group;

R is hydrogen or an isopropyl group; and

B is a heterocyclic base.

Treatment with an acid replaces the hydroxyl protecting group at the unbound terminus of the oligonucleotide, and thus enables the solid support bound oligomer to participate in the next synthetic iteration. This process is repeated until an oligomer of desired length is produced.

As will be appreciated, one or more of the 2'-, 3'-, and/or 5'-positions of the oligonucleotide comprises a hydroxyl protecting group. A wide variety of hydroxyl protecting groups can be employed in the methods of the invention. Preferably, the protecting group is stable under basic conditions but can be removed under acidic conditions. In general, protecting groups render chemical functionalities inert to specific reaction conditions, and can be appended to and removed from such functionalities in a molecule without substantially damaging the remainder of the molecule. Representative hydroxyl protecting groups are disclosed by Beaucage, *et al.*, *Tetrahedron* **1992**, 48, 2223-2311, and also in Greene and Wuts, *Protective Groups in Organic Synthesis*, Chapter 2, 2d ed, John Wiley & Sons, New York, 1991, each of which are hereby incorporated by reference in their entirety. Preferred protecting groups include

- 5 trimethoxytrityl, dimethoxytrityl (DMT), monomethoxytrityl, 9-phenylxanthen-9-yl (Pixyl) and 9-(p-methoxyphenyl)xanthen-9-yl (Mox).

The protecting group can be removed from oligomeric compounds of the invention by techniques well known in the art to form the free hydroxyl. A wide variety of bases can be used to initiate the removal of protecting groups. These bases include aqueous ammonium hydroxide, aqueous methylamine, DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) and carbonates containing counterions such as lithium, potassium, sodium, and cesium. Most preferred is potassium carbonate and ammonia. Removal of the protecting groups may be performed in a variety of suitable solvents. These solvents include those known to be suitable for protecting group removal in oligonucleotide synthesis. In the case of ammonia, water is the preferred solvent, whereas when using carbonates, alcohols are preferred. Methanol is most preferred. For example, dimethoxytrityl protecting groups can be removed by protic acids such as formic acid, dichloroacetic acid, trichloroacetic acid, p-toluene sulphonic acid or with Lewis acids such as for example zinc bromide. *See* for example, Greene and Wuts, *supra*. In addition, treatment of the oligonucleotide with ammonium hydroxide at room temperature for about 48 hours can be used to remove isopropyl substituted benzamidoethyl and cyanoethoxy protecting groups. The NH analogs of benzamidoethyl protecting groups, however, can be removed by treating the oligonucleotide with ammonium hydroxide at about 55°C for about 48 hours. Further, allyl protecting groups require the use of palladium zero and silyl protecting groups can be removed by fluoro. In certain preferred embodiments, conditions for removal of the oxygen or sulfur protecting group also effect cleavage of the oligomeric compound from the solid support. The deprotected terminal nucleotide and/or internucleosidic nucleotides can then be functionalized.

In some embodiments, phosphite or thiophosphite compounds are oxidized or sulfurized. The choice of oxidizing or sulfurizing agent will determine whether the linkage will be oxidized or sulfurized to a phosphotriester, thiophosphotriester, or a dithiophosphotriester linkage. Sulfurizing agents used during oxidation to form phosphorothioate and phosphorodithioate linkages include Beaucage reagent (*see e.g.* Iyer, R.P., *et. al.*, *J. Chem. Soc.*, **1990**, 112, 1253-1254, and Iyer, R.P., *et. al.*, *J. Org. Chem.*, **1990**, 55, 4693-4699); tetraethylthiuram disulfide



5 (see e.g., Vu, H., Hirschbein, B.L., *Tetrahedron Lett.*, **1991**, 32, 3005-3008); dibenzoyl  
tetrasulfide (see e.g., Rao, M.V., et. al., *Tetrahedron Lett.*, **1992**, 33, 4839-4842);  
di(phenylacetyl) disulfide (see e.g., Kamer, P.C.J., *Tetrahedron Lett.*, **1989**, 30, 6757-6760);  
Bis(O,O-diisopropoxy phosphinothioyl)disulfids (see Stec et al., *Tetrahedron Lett.*, **1993**, 34,  
5317-5320); 3-ethoxy-1,2,4-dithiazoline-5-one (see *Nucleic Acids Research*, **1996** 24, 1602-  
10 1607, and *Nucleic Acids Research*, **1996** 24, 3643-3644); Bis(p-chlorobenzenesulfonyl)disulfide  
(see *Nucleic Acids Research*, **1995** 23, 4029-4033); sulfur, sulfur in combination with ligands  
like triaryl, trialkyl, triaralkyl, or trialkaryl phosphines. The foregoing references are hereby  
incorporated by reference in their entirety.

Suitable oxidizing agents for forming the phosphodiester or phosphorothioate linkages  
15 include iodine/tetrahydrofuran/ water/pyridine or hydrogen peroxide/water or *tert*-butyl  
hydroperoxide or any peracid like *m*-chloroperbenzoic acid. In the case of sulfurization the  
reaction is performed under anhydrous conditions with the exclusion of air, in particular oxygen  
whereas in the case of oxidation the reaction can be performed under aqueous conditions.

As used herein, the terms "phosphorus protecting group" and "phosphorus blocking group"  
20 refers to a group that is initially bound to the phosphorus atom of a phosphoramidite. The  
phosphorus blocking group functions to protect the phosphorus containing internucleotide  
linkage or linkages during, for example, solid phase oligonucleotide synthetic regimes.  
Treatment of the internucleotide linkage or linkages that have a phosphorus blocking group  
thereon with a deblocking agent, such as aqueous ammonium hydroxide, will result in the  
25 removal of the phosphorus blocking group and leave a hydroxyl or thiol group in its place.

There are many phosphorus blocking groups known in the art which are useful in the  
present invention including, but not limited, to, cyanoethyl, diphenylsilylethyl, cyanobutenyl,  
cyano *p*-xylyl (CPX), methyl-N-trifluoroacetyl ethyl (META) and acetoxo phenoxy ethyl  
(APOE) groups. Phosphorus protecting groups are further described in Beaucage, S.L. and Iyer,  
30 R.P., *Tetrahedron*, **1993**, 49, 1925-1963; Beaucage, S.L. and Iyer, R.P., *Tetrahedron*, **1993**, 49,  
10441-10488; and Beaucage, S.L. and Iyer, R.P., *Tetrahedron*, **1992**, 48, 2223-2311.

Representative United States patents that teach the preparation of phosphorus protecting groups

5 and their incorporation into phosphoramidite compounds include, but are not limited to, United States Patent Nos. 5,783,690; 5,760,209; 5,705,621; 5,614,621; 5,453,496; 5,153,319; 5,132,418; 4,973,679; 4,725,677; 4,668,777; 4,500,707; 4,458,066; 4,415,732; and Re. 34,069, the entire contents of each of which are herein incorporated by reference.

10 As will be recognized, the steps of certain processes of the present invention need not be performed any particular number of times or in any particular sequence. Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following synthetic teachings and working examples which are intended to be illustrative of the present invention, and not limiting thereof.

## 15 EXAMPLES

As described in detail below, for the introduction of a labeled 5'-terminal phosphate or thiophosphate, phosphoramidites, **2** and **3** bearing pyrene or fluorescein reporter groups, respectively, were synthesized (Scheme 1, below). Similarly, phosphoramidite **4** (reference sample) and phosphoramidites **5-7** for 5'-terminal phosphorylation were obtained. The phosphoramidites **2-6** were incorporated in a standard manner at the last step of oligonucleotide synthesis (Scheme 3). The solid support bound oligonucleotides were deprotected with concentrated ammonium hydroxide to give crude **8-15**. On isolation, these were detritylated with diluted AcOH to give **16-23**. Brief treatment with a base removed the 5'-terminal 3-hydroxy-2,2-bis(ethoxycarbonyl) propyl-1 phosphate protecting group to afford **24-31**. The final products, 5'-phosphorylated **24** and **25** and 5'-labeled **26-31** were desalted by HPLC. All intermediates and final products were characterized by ESMS and HPLC (Tables 1 and 2, below).

For the introduction of 3'-terminal phosphate or thiophosphate, solid support **32** was synthesized (Scheme 2, below). The utility of **32** was verified in preparation of oligonucleotides **33-38** (Scheme 4 and Table 3, below). Oligonucleotide **32** demonstrated favorable properties in two respects. First, the TMT protecting group in **32** was removed with 3%  $\text{Cl}_2\text{HCCO}_2\text{H}$  in  $\text{CH}_2\text{Cl}_2$ , i.e. under milder conditions than the previously used for the removal of DMT group (3%

5  $F_2CCO_2H$  in  $CH_2Cl_2$ ). Second, the diglycolyl linker in **32** was cleaved with aqueous bases faster than previously reported for the malonyl linker.

The reactivity of the 3'-terminal phosphorothioate group was verified by labeling **37** with *N*-(1-pyrenylmethyl)iodoacetamide and 5-iodoacetamidofluorescein to afford **39** and **40** (Scheme 5 and Table 3, below).

10 The preparation of oligonucleotides labeled with two different fluorescent reporter groups at the 5'- and 3'-termini was accomplished according to Schemes 6 and 7. First, protected oligonucleotides **41a** and **41b** were assembled on solid support **32**. For the last coupling, 5'-*O*-(4,4'-dimethoxytrityl) - 3'-*O*-(*N,N*- diisopropylamino) [2-(4- methoxybenzamido) ethoxy] phosphinyl-2'-deoxythymidine was used. For the first and the last coupling cycles,  
15 sulfurization with 3*H*-1,2-benzodithiol-3-one 1,1-dioxide was carried out while all other cycles employed oxidation with *t*-BuOOH in MeCN. Thus, the oligonucleotides synthesized contained the 3'-terminal phosphorothioate and a 2-(4-methoxybenzamido)ethyl protected P=S linkage preceding the 5'-terminal nucleoside residue. After deprotection with ammonium hydroxide (2 days/RT), **42** was obtained and isolated by HPLC (Table 4). On detritylation, **42** gave **43**, which  
20 was reacted with **44** to give **45**. Deprotection of **45** with concentrated ammonium hydroxide for 48 h at 55 °C removed 2-(4-methoxybenzamido)ethyl protection to give **46** which contained the second phosphorothioate group for the conjugation. Compound **46** was next labeled with **44** and **47-49** to give a bis-pyrenyl labeled **50** and unsymmetrically labeled **51-53** (Scheme 7 and Table 4). The methodology described above allows one to place the second reporter group in any  
25 internucleosidic position of the oligonucleotide.


The methodology described above was also used to label 5'- and 3'-terminal phosphorothioate residues with two different reporter groups. First, solid support-bound oligonucleotides **54a** and **b** bearing the selectively protected 5'-terminal phosphorothioate group were synthesized on solid support **32** as depicted in Scheme 8. These were treated with  
30 ammonium hydroxide for 2 days at RT to give **55** where only the 3'-terminal phosphorothioate group was deprotected. Compound **55** was next labeled with **49**. The product was isolated by HPLC, and 2-(4-methoxybenzamido)ethyl protection was removed with concentrated

ammonium hydroxide for 48 h at 55 °C. Without isolation, the product was detritylated and briefly treated with a base, which removed diethyl 3-hydroxy-*bis*-(ethoxycarbonyl)propyl protection from the 5'-terminal phosphorothioate group to give **56**. This was treated with labeling reagents **44**, **48**, and **49**, and the unsymmetrically bis-labeled oligonucleotides **57-59** were isolated by HPLC and characterized (Scheme 9 and Table 5, below).

EXAMPLE 1. *Diethyl 2-[(4,4',4''-trimethoxytrityl)oxymethyl]-2-hydroxymethylmalonate (1)*.

Referring to scheme 1, diethyl *bis*-(hydroxymethyl)malonate (11.5 g, 55 mmol) was treated overnight with 4,4',4''-trimethoxytrityl chloride (19.2 g, 52 mmol) in pyridine (16 mL) and dioxane (100 mL), and the solvent was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (500 mL), washed with 5% aqueous NaHCO<sub>3</sub> (3 × 100 mL), washed with brine (2 × 100 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The residue was purified on a silica gel column using a step gradient of ethyl acetate (0 to 15%) in toluene to give **1** (8.2 g, 67.5%) as an oil. HR MALDI MS: calculated for C<sub>31</sub>H<sub>36</sub>O<sub>9</sub> (M + H<sup>+</sup>) 553.2442, found 553.2438. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.29 (6H, d, *J* = 8.8 Hz), 6.83 (6H, d, *J* = 8.8 Hz), 4.26-4.11 (6H, m), 3.80 (9H, s), 3.62 (2H, s), 2.11 (1H, t, *J* = 5.8 Hz), 1.24 (6H, t, *J* = 7.0 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 169.3 (C), 158.5 (C), 136.1 (C), 129.8 (CH), 113.2 (CH), 86.0 (C), 63.7 (CH<sub>2</sub>), 61.9 (CH<sub>2</sub>), 61.6 (CH<sub>2</sub>), 60.6 (C), 55.2 (CH<sub>3</sub>), 14.1 (CH<sub>3</sub>).

Scheme 1



EXAMPLE 2. Diethyl [[[tris (4 - methoxyphenyl) methyl] oxy] methyl] [[[[bis (1 - methylethyl) amino] [6 - [[3',6' - bis (2,2 - dimethylpropionyloxy) - 3 - oxospiro [isobenzofuran - 1 (3H),9' - [9H] xanthen] - 5 - yl] carboxamido] hexyloxy] phosphino] oxy] methyl] propanedioate (3).

A solution of chloro bis[(N,N-diisopropyl)amino]phosphite (768 mg, 2.88 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added dropwise to a mixture of 1 (1381 mg, 2.5 mmol) and N-ethyl-N,N-diisopropylamine (485 mg, 3.75 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL) under magnetic stirring at -30°C. The reaction mixture was allowed to warm up to room temperature, and the stirring was continued for 1 h. 6-[[3',6'-bi (2,2-dimethylpropionyloxy)-3- oxospiro[isobenzofuran-1(3H),9'-[9H]xanthen]-5-yl] carboxamido]hexanol (1030 mg, 1.6 mmol) was added followed

5 by 1*H*-tetrazole (0.45 M in MeCN; 2.8 mL, 1.25 mmol). The resulting mixture was kept at room temperature for 2 h. Aqueous NaHCO<sub>3</sub> (5%; 5 mL) was added, the emulsion was diluted with brine (25 mL), and the product was extracted with ethyl acetate (3 × 50 mL). Extracts were washed with brine (3 × 25 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was dissolved in toluene (50 mL), applied on a silica gel column, and separated eluting with a gradient from  
10 5:91:4 to 40:56:4 ethyl acetate / hexane / triethylamine. Collected fractions were evaporated, co-evaporated with dry MeCN (3 × 50 mL), dry toluene (3 × 50 mL), and dried on an oil pump to give **3** (1400 mg, 66.0%) as a yellow foam. HR FAB MS: found *m/z* 1324.5850; C<sub>74</sub>H<sub>89</sub>N<sub>2</sub>O<sub>18</sub>P requires 1324.5848. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.06 (1H, dd, *J* = 1.3, 7.9 Hz), 8.04 (1H, dd, *J* = 1.3, 7.9 Hz), 7.57 (1H, dd, *J* = 1.3 and 1.4 Hz), 7.25-7.19 (6H, m), 7.11 (1H, d, *J* = 1.4 Hz), 7.10 (1H, d, *J* = 1.4 Hz), 7.02 (1H, t, *J* = 5.8 Hz), 6.93-6.89 (2H, m), 6.84 (1H, dd, *J* = 0.9 and 2.3 Hz), 6.82-6.78 (7H, m), 4.22 (1H, dd, *J* = 5.7 and 9.7 Hz), 4.16-3.97 (7H, m), 3.71 (9H, s), 3.54 (1H, d, *J* = 8.5 Hz), 3.51 (1H, d, *J* = 8.5 Hz), 3.52-3.41 (2H, m), 3.23-3.16 (2H, m), 1.50-1.36 (4H, m), 1.32 (18H, s), 1.32-1.20 (4H, m), 1.12 (3H, t, *J* = 7.1 Hz), 1.11 (3H, t, *J* = 7.0 Hz), 1.08 (6H, d, *J* = 6.8 Hz), 1.03 (6H, d, *J* = 6.6 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 177.5 (C), 169.2 (C), 169.12 (C), 169.09 (C),  
20 166.1 (C), 159.5 (C), 154.0 (C), 152.4 (C), 143.0 (C), 137.2 (C), 130.7 (CH), 130.4 (CH), 130.3 (CH), 128.8 (C), 126.2 (CH), 123.4 (CH), 119.2 (CH), 118.3 (CH), 117.0 (C), 113.9 (CH), 111.4 (CH), 86.5 (C), 82.5 (C), 64.0 (CH<sub>2</sub>), 62.6 (CH<sub>2</sub>), 62.3 (CH<sub>2</sub>), 62.2 (CH<sub>2</sub>), 61.7 (CH<sub>2</sub>), 60.7 (C), 55.8 (CH<sub>3</sub>), 43.7 (CH), 43.6 (CH), 40.7 (CH<sub>2</sub>), 39.8 (C), 31.8 (CH<sub>2</sub>), 29.9 (CH<sub>2</sub>), 27.3 (CH<sub>2</sub>), 27.2 (CH<sub>3</sub>), 26.4 (CH<sub>2</sub>), 25.0 (CH<sub>3</sub>), 24.9 (CH<sub>3</sub>), 24.8 (CH<sub>3</sub>), 14.3 (CH<sub>3</sub>). <sup>31</sup>P NMR (CDCl<sub>3</sub>) δ 147.4.

25

EXAMPLE 3. Diethyl [[[bis (4-methoxyphenyl)phenylmethyl]oxy] methyl][[[bis (1-methylethyl) amino] [6 - [[3',6'- bis (2,2-dimethylpropionyloxy) - 3 - oxospiro [isobenzofuran - 1 (3*H*), 9' - [9*H*] xanthen] - 5 - yl] carboxamido] hexyloxy] phosphino] oxy] methyl] propanedioate (**3a**).

30

Compound **3a** was prepared analogously from diethyl 2-[[[bis (4-methoxyphenyl) phenylmethyl] oxy] methyl] -2-hydroxymethylpropanedioate **1a** (829 mg, 1.5 mmol), chloro bis[(*N,N*-diisopropyl)amino]phosphite (440 mg, 1.65 mmol), and 6-[[3',6'-bi (2,2-

5 dimethylpropionyloxy)-3-oxospiro[isobenzofuran-1(3H),9'-[9H] xanthen]-5-yl]  
carboxamido]hexanol (483 mg, 0.5 mmol). Isolation on a silica gel column using gradient from  
5:93:2 to 40:58:2 ethyl acetate / hexane / triethylamine gave **3a** (798 mg, 82.1%) as a yellow  
foam. HR FAB MS: found  $m/z$  1294.5742;  $C_{73}H_{87}N_2O_{17}P$  requires 1294.5742.  $^1H$  NMR ( $CDCl_3$ )  
8 8.10 (1H, dd,  $J = 1.3$  and 7.9 Hz), 8.02 (1H, dd,  $J = 1.3$  and 7.9 Hz), 7.54 (1H, dd,  $J = 1.3$  and  
10 1.4 Hz), 7.45-7.15 (9H, m), 7.11 (1H, d,  $J = 1.4$  Hz), 7.08 (1H, d,  $J = 1.4$  Hz), 7.02 (1H, t,  $J = 5.8$   
Hz), 6.92-6.87 (2H, m), 6.82-6.78 (6H, m), 4.20 (1H, dd,  $J = 5.7$  and 9.7 Hz), 4.16-3.95 (7H, m),  
3.78 (6H, s), 3.58 (1H, d,  $J = 8.5$  Hz), 3.50 (1H, d,  $J = 8.5$  Hz), 3.51-3.41 (2H, m), 3.24-3.16 (2H,  
m), 1.52-1.36 (4H, m), 1.34 (18H, s), 1.32-1.20 (4H, m), 1.12 (3H, t,  $J = 7.0$  Hz), 1.11 (3H, t,  $J =$   
6.9 Hz), 1.07 (6H, d,  $J = 6.7$  Hz), 1.00 (6H, d,  $J = 6.7$  Hz).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  176.6 (C), 168.5  
15 (C), 168.4 (C), 165.5 (C), 158.4 (C), 153.4 (C), 152.8 (C), 151.5 (C), 144.8 (C), 141.6 (C), 135.9  
(C), 130.2 (CH), 129.3 (CH), 129.0 (CH), 128.3 (CH), 127.7 (CH), 126.7 (CH), 125.5 (CH),  
122.4 (CH), 117.9 (CH), 115.6 (C), 113.0 (CH), 110.5 (CH), 85.8 (C), 81.9 (C), 63.3, 63.0  
(CH<sub>2</sub>), 62.0, 61.7 (CH<sub>2</sub>), 61.2 (CH<sub>2</sub>), 60.9 (CH<sub>2</sub>), 60.0, 59.9 (C), 55.2 (CH<sub>3</sub>), 43.0, 42.8 (CH),  
40.4 (CH<sub>2</sub>), 39.2 (CH<sub>2</sub>), 31.1, 31.0 (CH<sub>2</sub>), 29.3 (CH<sub>2</sub>), 27.1 (CH<sub>3</sub>), 26.6 (CH<sub>2</sub>), 25.6 (CH<sub>2</sub>), 24.6,  
20 24.5 (CH<sub>3</sub>), 14.1 (CH<sub>3</sub>).  $^{31}P$  NMR ( $CDCl_3$ )  $\delta$  146.9.

EXAMPLE 4. Diethyl [[[tris (4-methoxyphenyl) methyl] oxy] methyl] [[[[bis (1-methylethyl)  
amino] [4-(1-pyrenebutyl)oxy] phosphino] oxy] methyl] propanedioate (**2**).

Compound **2** was prepared from **1** (1630 mg, 2.95 mmol), chloro bis[(*N,N*,-  
25 diisopropyl)amino]phosphite (944 mg, 3.54 mmol), and 4-(1-pyrene)butanol (1015 mg, 3.7  
mmol) as described for **3**. Isolation on a silica gel column using gradient from 5:91:4 to 40:56:4  
ethyl acetate / hexane / triethylamine gave **2** (2185 mg, 77.5%) as a white foam. HR FAB MS:  
found  $m/z$  955.4421;  $C_{57}H_{67}NO_{10}P$  requires 955.4424.  $^1H$  NMR ( $CDCl_3$ )  $\delta$  8.24 (1H, d,  $J = 9.2$   
Hz), 8.12 (2H, d,  $J = 7.5$  Hz), 8.07-7.93 (5H, m), 7.80 (1H, d,  $J = 7.9$  Hz), 7.31-7.25 (6H, m),  
30 6.78-6.72 (6H, m), 4.37 (1H, dd,  $J = 5.8$  and 9.8 Hz), 4.26 (1H, dd,  $J = 3.8$  and 9.8 Hz), 4.21-  
4.01 (4H, m), 3.71 (2H, s), 3.74-3.43 (4H, m), 3.66 (9H, s), 3.30 (2H, t,  $J = 7.7$  Hz), 2.0-1.86  
(2H, m), 1.75-1.68 (2H, m), 1.14 (3H, t,  $J = 7.2$  Hz), 1.13 (3H, t,  $J = 7.0$  Hz), 1.11 (6H, d,  $J = 6.4$

5 Hz), 1.07 (6H, d,  $J = 6.6$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  168.4 (C), 168.3 (C), 158.2 (C), 137.0 (C), 136.4 (C), 131.4 (C), 130.9 (C), 129.9 (CH), 128.6 (C), 127.5 (CH), 127.2 (CH), 127.1 (CH), 126.4 (CH), 125.7 (CH), 125.1 (C), 125.0 (CH), 124.8 (CH), 124.7 (CH), 124.6 (CH), 123.5 (CH), 112.9 (CH), 85.5 (C), 63.2 ( $\text{CH}_2$ ), 61.8 ( $\text{CH}_2$ ), 61.1 ( $\text{CH}_2$ ), 60.9 ( $\text{CH}_2$ ), 60.0 ( $\text{CH}_2$ ), 59.9 (C), 55.0 ( $\text{CH}_3$ ), 43.0 (CH), 42.8 (CH), 33.1 ( $\text{CH}_2$ ), 31.3 ( $\text{CH}_2$ ), 28.2 ( $\text{CH}_2$ ), 24.5 ( $\text{CH}_3$ ), 24.4 (CH<sub>3</sub>), 13.9 (CH<sub>3</sub>).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  150.0.

EXAMPLE 5. Diethyl [[[tris (4-methoxyphenyl) methyl] oxy] methyl] [[[[bis (1-methylethyl) amino] (ethoxy) phosphino] oxy] methyl] propanedioate (4).

Compound 4 was prepared from 1 (553 mg, 1.0 mmol), chloro bis[(*N,N*,  
15 diisopropyl)amino]phosphite (307 mg, 1.15 mmol), and ethanol (69 mg, 1.5 mmol) as described for 3. Isolation on a silica gel column using gradient from 5:93:2 to 40:58:2 ethyl acetate / hexane / triethylamine gave 4 (662 mg, 91%) as a colorless oil. HR FAB MS: found  $m/z$  727.3486;  $\text{C}_{39}\text{H}_{54}\text{NO}_{10}\text{P}$  requires 727.3485.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.40-7.27 (6H, m), 6.92-6.67 (6H, m), 4.37 (1H, dd,  $J = 5.8$  and 7.5 Hz), 4.26-4.02 (5H, m), 3.77 (9H, s), 3.68 (2H, s), 3.75-  
20 3.39 (2H, m), 3.20 (2H, m), 1.40-1.00 (21H, m).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  168.6 (C), 168.4 (C), 158.3 (C), 136.5 (C), 130.0 (CH), 113.0 (CH), 85.6 (C), 63.7 ( $\text{CH}_2$ ), 61.6 ( $\text{CH}_2$ ), 61.3 ( $\text{CH}_2$ ), 60.9 ( $\text{CH}_2$ ), 60.0 ( $\text{CH}_2$ ), 58.9 (C), 55.2 ( $\text{CH}_3$ ), 43.0 (CH), 42.8 (CH), 24.6 ( $\text{CH}_3$ ), 24.4 ( $\text{CH}_3$ ), 19.1 ( $\text{CH}_3$ ), 14.0 ( $\text{CH}_3$ ).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  145.6.

25 EXAMPLE . Diethyl [[[tris (4-methoxyphenyl) methyl] oxy] methyl] [[[[bis (1-methylethyl) amino] [2-(4-methoxybenzamido)ethoxy] phosphino] oxy] methyl] propanedioate (5).

Compound 5 was prepared from 1 (553 mg, 1.0 mmol), chloro bis[(*N,N*,  
diisopropyl)amino]phosphite (307 mg, 1.15 mmol), and 2-(4-methoxybenzamido)ethanol (254  
mg, 1.3 mmol) as described for 3. Isolation on a silica gel column using gradient from 0:95:5 to  
30 40:55:5 ethyl acetate / hexane / triethylamine gave 5 (652 mg, 74.4%) as a colorless oil. HR FAB MS: found  $m/z$  876.3963;  $\text{C}_{47}\text{H}_{61}\text{N}_2\text{O}_{12}\text{P}$  requires 876.3962.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.8-7.65 (2H, m); 7.40-7.27 (7H, m), 6.92-6.67 (8H, m), 4.3-4.0 (6H, m), 3.81 (3H, s); 3.77 (9H, s), 3.7-3.3 (8H,



5 m), 1.40-1.00 (18H, m).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  168.6 (C), 168.4 (C), 168.1 (C), 162.1 (C), 158.0 (C), 136.3 (C), 130.2 (CH), 128.8 (CH), 126.4 (C), 117.5 (C), 113.7 (CH), 113.1 (CH), 85.6 (C), 63.7, 63.4 ( $\text{CH}_2$ ), 62.6, 62.2 ( $\text{CH}_2$ ), 62.1, 61.9 ( $\text{CH}_2$ ), 61.4 ( $\text{CH}_2$ ), 61.3 (C), 55.3 ( $\text{CH}_3$ ), 55.1 ( $\text{CH}_3$ ), 43.1 (CH), 42.9 (CH), 40.9 ( $\text{CH}_2$ ), 24.6 ( $\text{CH}_3$ ), 24.4 ( $\text{CH}_3$ ), 14.2 ( $\text{CH}_3$ ).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  147.9.

10  
EXAMPLE 6. Diethyl [[[tris (4 – methoxyphenyl) methyl] oxy] methyl] [[[[bis (1 – methylethyl) amino] (2–cyanoethoxy) phosphino] oxy] methyl] propanedioate (6).

1H–Tetrazole (0.5 mmol, 1.1 mL of 0.45 M solution in MeCN) was added to 1 (553 mg, 1.0 mmol) and 2–cyanoethyl *N,N,N',N'*-tetraisopropyl phosphorodiamidite (332 mg, 1.1 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL). The solution was stirred for 2 h, and 5% aqueous  $\text{NaHCO}_3$  (5 mL) was added. The mixture was diluted with brine (15 mL) and the product was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 40$  mL). The extracts were dried over  $\text{Na}_2\text{SO}_4$  and evaporated. The residue was purified on a silica gel column eluting with a gradient from 0:95:5 to 30:65:5 ethyl acetate / hexane / triethylamine to give 6 (650 mg, 86.3%) as a colorless oil. HR FAB MS: found  $m/z$  752.8309;  $\text{C}_{40}\text{H}_{53}\text{N}_2\text{O}_{10}\text{P}$  requires 752.8301.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.40-7.25 (6H, m), 6.9-6.65 (6H, m), 4.3-4.0 (6H, m), 3.77 (9H, s), 3.65 (2H, s), 3.8-3.4 (4H, m), 2.55 (2H, m), 1.40-1.00 (18H, m).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  168.6 (C), 168.4 (C), 158.3 (C), 136.5 (C), 130.0 (CH), 117.3 (C), 113.0 (CH), 85.6 (C), 63.9, 63.6 ( $\text{CH}_2$ ), 62.1, 61.9 ( $\text{CH}_2$ ), 61.4 ( $\text{CH}_2$ ), 61.3 (C), 61.1, 60.9 ( $\text{CH}_2$ ), 55.1 ( $\text{CH}_3$ ), 43.1 (CH), 42.9 (CH), 24.6 ( $\text{CH}_3$ ), 24.4 ( $\text{CH}_3$ ), 20.5 ( $\text{CH}_2$ ), 14.2 ( $\text{CH}_3$ ).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  147.4.

25  
EXAMPLE 7. Diethyl [[[tris (4 – methoxyphenyl) methyl] oxy] methyl] [[[[bis (1 – methylethyl) amino] [(9H–fluorene–9–yl)methoxy] phosphino] oxy] methyl] propanedioate (7).

Compound 7 was prepared from 1 (1430 mg, 2.6 mmol), chloro bis[(*N,N*,-diisopropyl)amino]phosphite (827 mg, 3.1 mmol), and (9H–fluorene–9–yl)methanol (712 mg, 3.6 mmol) as described for 3. Isolation on a silica gel column using gradient from 0:99:1 to 25:74:1 ethyl acetate / hexane / triethylamine gave 7 (1548 mg, 68.1%) as a white solid foam. HR FAB MS: found  $m/z$  727.3486;  $\text{C}_{39}\text{H}_{54}\text{NO}_{10}\text{P}$  requires 727.3485.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.8-7.5


5 (4H, m); 7.5-7.15 (10H, m); 6.9-6.7 (6H, m); 4.50-4.22 (2H, m); 4.20-4.05 (4H, m); 4.05-3.85 (2H, m); 3.72 (9H, s); 3.80-3.40 (5H, m); 1.30-1.00 (18H, m).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  147.4.

EXAMPLE 8. *Preparation of the solid support 32.*

Referring to scheme 2, a mixture of **1** (1.11 g, 2.0 mmol), 1,4-dioxane-2,6-dione (0.70 g, 6.0 mmol), Py (5 mL), and dioxane (5 mL) was kept overnight at room temperature. The solvent was  
10 evaporated and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with 2 M aqueous triethylammonium acetate ( $5 \times 10$  mL) and water ( $5 \times 10$  mL), dried over  $\text{Na}_2\text{SO}_4$ . The extract was evaporated to give crude monoester of **2** with diglycolic acid (1.54 g; triethylammonium salt) as a solid foam in a quantitative yield.

15

Scheme 2



Crude product from the previous step (0.99 g, 1.28 mmol) and triphenylphosphine (0.40 g, 1.5 mmol) were dissolved in 1,2-dichloroethane (5 mL). To this was added a solution of 3,3'-  
20 dinitrodipyridyl disulfide (0.47 g, 1.5 mmol) and DMAP (0.18 g, 1.5 mmol) in 1,2-dichloroethane (5 mL). The mixture was shaken for 15 min and filtered. The precipitate was washed on filter with 1,2-dichloroethane (5 mL), and the combined filtrates were added to aminoalkyl CPG (5.98 g;  $119 \mu\text{mol g}^{-1}$ , 0.71 mmol). The suspension was shaken for 1 h and filtered. The solid support was washed with 1,2-dichloroethane ( $3 \times 10$  mL), treated with  
25  $\text{Ac}_2\text{O}/\text{N}$ -methylimidazole/THF (10:10:80) for 30 min, washed with 1,2-dichloroethane ( $5 \times 10$  mL) and MeCN ( $5 \times 10$  mL), and dried to give **32**. An aliquot of **32** was treated with TFA (2% in








- 5  $\text{CH}_2\text{Cl}_2$ ), and the concentration of the released trimethoxytrityl cation was determined colorimetrically at 486 nm to give the loading of  $80.5 \mu\text{mol g}^{-1}$ .

### EXAMPLE 9. Oligonucleotide synthesis.

- Referring to scheme 3, oligonucleotide synthesis was carried out on an ABI 380B DNA  
 10 Synthesizer on 1 to 20  $\mu\text{mol}$  scale using phosphoramidite chemistry. For the coupling step, 0.1 M solutions of phosphoramidite building blocks in MeCN (for 2, 0.1 M solution in  $\text{CH}_2\text{Cl}_2$ :MeCN 50:50, v/v or in  $\text{CH}_2\text{Cl}_2$ ) were used; 0.45 M 1H-tetrazole was used as an activator. The coupling time for phosphoramidites 2, 3, and 5 was 600 s.

15

Scheme 3



5

All oligonucleotides were assembled using the standard base protection strategy: *N*-benzoyl protected dA, dC, 2'-*O*-(2-methoxyethyl)-A, and 2'-*O*-(2-methoxyethyl)-5-methyl-C phosphoramidites and *N*-isobutyryl protected dG and 2'-*O*-(2-methoxyethyl)-G phosphoramidites. Additionally, 9-12 were assembled using *N*-phenoxyacetyl dA, dC, and dG phosphoramidites. For the preparation of oligonucleotides 8-13, 33-35, 41b, and 54b, phosphoramidites protected with 2-cyanoethyl group at the P(III) were used.<sup>5</sup> For the synthesis of oligonucleotides 41a and 54a phosphoramidites protected with *N*-isopropyl-(4-methoxybenzamido)ethyl group were used.<sup>1</sup> The 5'-terminal phosphorylation in preparation of 8 and 9 was performed using phosphoramidites 6 and 7 to demonstrate a very similar performance of both reagents. For preparation of 41a and 41b, the last coupling was carried out using 5'-*O*-(4,4'-dimethoxytrityl)-3'-*O*-(*N,N*-diisopropylamino)[2-(4-methoxybenzamido)ethoxy]phosphinyl-2'-deoxythymidine as a building block.<sup>1</sup>

For the oxidation step, a commercial iodine oxidizer or *t*-BuOOH (10% in MeCN) were used. For sulfurization, 3*H*-1,2-benzodithiol-3-one 1,1-dioxide<sup>4</sup> (0.05 M in MeCN) was used as the sulfur-transfer reagent. Thus, oligonucleotides 8, 11, 14, 15, 33 were assembled using the iodine oxidizer. Oligonucleotides 9, 10, 12, 13, 35 were assembled using 3*H*-1,2-benzodithiol-3-one 1,1-dioxide. Oligonucleotides 34, 41a, 41b, 54a, and 54b that contained both P=S and P=O linkages were assembled using 3*H*-1,2-benzodithiol-3-one 1,1-dioxide for sulfuration and *t*-BuOOH for oxidation.


25

EXAMPLE 10. *Releasing oligonucleotides from solid support and deprotection.*

Referring to scheme 4, for all oligonucleotides, concentrated ammonium hydroxide was used as a deprotecting agent. On completing the chain assembly, the oligonucleotides 8, 14, and 15 were treated for 2 h at room temperature. The oligonucleotides 9, 33-35, and 41 were deprotected for 6 h at 55 °C.

30

Scheme 4



Under optimized conditions, the oligonucleotides **10–13** and **54** that possessed 5'-terminal phosphotriester moiety and phenoxyacetyl as the base protecting groups were treated at room temperature for 2 h. Alternatively, when nucleic bases were protected with the standard protecting groups, **10–13** and **54** were deprotected for 2 days at room temperature. On evaporation, the deprotected oligonucleotides were isolated by reversed phase HPLC and characterized by ESMS as described below.

#### EXAMPLE 11. *Detritylation procedure.*

Oligonucleotides **8**, **9**, **33–35**, and **42** were deprotected with 10% aqueous AcOH (3 mL and 100 mL for 1 and 20  $\mu\text{mol}$  scale, correspondingly) for 30 min.

Oligonucleotides **10–15**, **25** were dissolved in 10% aqueous AcOH (3 mL and 100 mL for 1 and 20  $\mu\text{mol}$  scale, correspondingly). When the desalted material was subjected to the removal

- 5 of TMT protecting group, the deprotection was complete in 90 min. When the samples of oligonucleotides contained the HPLC buffer,  $\text{NH}_4\text{OAc}$ , the reaction mixture was kept for 4 to 5 h at room temperature.

When the detritylation was complete, the reaction mixtures were evaporated. The oligonucleotides **36-38** and **43** were desalted and characterized (Tables 3 and 4, respectively).

- 10 The oligonucleotides **16-22** (Tables 1 and 2) were coevaporated with water and the 5'-terminal 3-hydroxy-2,2-bis(ethoxycarbonyl)propyl-1 group was removed as described below.

5 Table 1. HPLC Retention Times for Oligonucleotides 8-31.

Compound	R	R <sup>1</sup>	Sequence, 5' to 3'	X <sup>a</sup>	Retention time, min <sup>b</sup>	
					Gradient 1	Gradient 2
8	TMT	H	T <sub>12</sub> <SEQ. ID. NO. 1>	O	32.8	—
9	TMT	H	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	27.7	23.8
10	TMT	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	—	30.2, 30.7
11	TMT	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	O	—	29.1
12	TMT	Flu	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	33.4, 34.2	—
13	TMT	Flu	AGCT <sub>2</sub> C T <sub>3</sub> GCACA TGTA <sub>3</sub> <sup>c</sup> <SEQ. ID. NO. 3>	S	35.9, 36.3	—
14	TMT	Flu	T <sub>12</sub> <SEQ. ID. NO. 1>	O	32.9	—
15	TMT	C <sub>2</sub> H <sub>5</sub>	T <sub>12</sub> <SEQ. ID. NO. 1>	O	33.0	—
16	H	H	T <sub>12</sub> <SEQ. ID. NO. 1>	O	18.0	—
17	H	H	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	19.6	16.4
18	H	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	—	25.0
19	H	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	O	—	24.5
20	H	Flu	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	23.4	—
21	H	Flu	AGCT <sub>2</sub> C T <sub>3</sub> GCACA TGTA <sub>3</sub> <sup>c</sup> <SEQ. ID. NO. 3>	S	28.5	—
22	H	Flu	T <sub>12</sub> <SEQ. ID. NO. 1>	O	22.7	—
23	H	C <sub>2</sub> H <sub>5</sub>	T <sub>12</sub> <SEQ. ID. NO. 1>	O	21.5	—
24	—	H	T <sub>12</sub> <SEQ. ID. NO. 1>	O	16.2	—
25	—	H	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	18.0	15.2
26	—	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	22.6	17.1
27	—	Pyr	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	O	22.0	16.7
28	—	Flu	ATGCAT <sub>2</sub> CTGC <sub>3</sub> A <sub>2</sub> G <sub>2</sub> A <SEQ. ID. NO. 2>	S	19.1	14.5
29	—	Flu	AGCT <sub>2</sub> C T <sub>3</sub> GCACA TGTA <sub>3</sub> <sup>c</sup> <SEQ. ID. NO. 3>	S	21.7	—
30	—	Flu	T <sub>12</sub> <SEQ. ID. NO. 1>	O	18.1	—
31	—	C <sub>2</sub> H <sub>5</sub>	T <sub>12</sub> <SEQ. ID. NO. 1>	O	17.9	—

<sup>a</sup> All oligonucleotides contained uniform, either phosphate (X=O) or phosphorothioate (X=S) backbone;<sup>b</sup> For HPLC conditions, consult Experimental Procedures; <sup>c</sup> 2'-O-(2-methoxyethyl) ribonucleotide residues are italicized; C stands for 5-methyl-2'-O-(2-methoxyethyl)cytidine residue.

5 Table 2.UV and ESMS data for Oligonucleotides 8–31.

Compound	Reporter group, $\lambda_{\max}$ , nm	ES MS		
		found, amu	Molecular Formula	calc., amu
8	—	4202.8	$C_{151}H_{192}N_{24}O_{93}P_{12}$	4202.9
9	—	6998.0	$C_{224}H_{280}N_{74}O_{108}P_{20}S_{20}$	6997.8
10	313, 327, 343	7252.8	$C_{244}H_{296}N_{74}O_{108}P_{20}S_{20}$	7254.2
11	314, 326, 343	6932.8	$C_{244}H_{296}N_{74}O_{128}P_{20}$	6932.9
12	463, 496	7455.5	$C_{251}H_{303}N_{75}O_{114}P_{20}S_{20}$	7455.3
13	463, 496	8402.4	$C_{292}H_{382}N_{72}O_{140}P_{20}S_{20}$	8401.4
14	463, 496	4660.7	$C_{178}H_{215}N_{25}O_{99}P_{12}$	4660.4
15	—	4229.2	$C_{153}H_{196}N_{24}O_{93}P_{12}$	4231.0
16	—	3870.5	$C_{129}H_{172}N_{24}O_{90}P_{12}$	3870.5
17	—	6665.5	$C_{202}H_{260}N_{74}O_{105}P_{20}S_{20}$	6665.4
18	322, 336, 352	6920.6	$C_{222}H_{276}N_{74}O_{105}P_{20}S_{20}$	6921.8
19	322, 335, 353	6599.1	$C_{222}H_{276}N_{74}O_{125}P_{20}$	6600.5
20	456, 472	7122.1	$C_{229}H_{283}N_{75}O_{111}P_{20}S_{20}$	7122.9
21	458, 482	8067.2	$C_{270}H_{362}N_{72}O_{137}P_{20}S_{20}$	8069.0
22	456, 472	4327.0	$C_{156}H_{195}N_{25}O_{96}P_{12}$	4328.0
23	—	3897.7	$C_{131}H_{176}N_{24}O_{90}P_{12}$	3898.6
24	—	3668.1	$C_{120}H_{158}N_{24}O_{85}P_{12}$	3668.3
25	—	6463.4	$C_{193}H_{246}N_{74}O_{100}P_{20}S_{20}$	6463.2
26	322, 336, 352	6720.1	$C_{213}H_{262}N_{74}O_{100}P_{20}S_{20}$	6719.6
27	322, 336, 352	6400.3	$C_{213}H_{262}N_{74}O_{120}P_{20}$	6398.3
28	458, 472	6919.6	$C_{220}H_{269}N_{75}O_{106}P_{20}S_{20}$	6920.7
29	456, 482	7869.0	$C_{261}H_{348}N_{72}O_{132}P_{20}S_{20}$	7866.8
30	456, 472	4125.4	$C_{147}H_{181}N_{25}O_{91}P_{12}$	4125.8
31	—	3695.8	$C_{122}H_{164}N_{24}O_{85}P_{12}$	3696.4



5 Table 3.ESMS data oligonucleotides 33-40.<sup>a</sup>

	Sequence, 5' to 3'	Backbone	ESMS, found	Molecular Formula	ESMS, calculated
33	DMTr-TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpO <SEQ. ID. NO. 4>	P=O	6365.2	C <sub>211</sub> H <sub>263</sub> N <sub>71</sub> O <sub>122</sub> P <sub>20</sub>	6365.2
34	DMTr-TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpS <SEQ. ID. NO. 5>	P=O	6381.7	C <sub>211</sub> H <sub>263</sub> N <sub>71</sub> O <sub>121</sub> P <sub>20</sub> S	6381.3
35	DMTr-TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpS <SEQ. ID. NO. 5>	P=S	6686.9	C <sub>211</sub> H <sub>263</sub> N <sub>71</sub> O <sub>102</sub> P <sub>20</sub> S <sub>20</sub>	6686.5
36	TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpO <SEQ. ID. NO. 6>	P=O	6062.2	C <sub>190</sub> H <sub>245</sub> N <sub>71</sub> O <sub>120</sub> P <sub>20</sub>	6062.8
37	TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpS <SEQ. ID. NO. 7>	P=O	6079.4	C <sub>190</sub> H <sub>245</sub> N <sub>71</sub> O <sub>119</sub> P <sub>20</sub> S	6078.9
38	TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> AtpS <SEQ. ID. NO. 7>	P=S	6384.7	C <sub>190</sub> H <sub>245</sub> N <sub>71</sub> O <sub>100</sub> P <sub>20</sub> S <sub>20</sub>	6384.2
39	TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> ATpS-Pyr <SEQ. ID. NO. 8>	P=O	6350.9	C <sub>209</sub> H <sub>258</sub> N <sub>72</sub> O <sub>120</sub> P <sub>20</sub> S	6350.2
40	TGCATC <sub>5</sub> AG <sub>2</sub> C <sub>2</sub> AC <sub>2</sub> ATpS-Flu <SEQ. ID. NO. 9>	P=O	6467.0	C <sub>212</sub> H <sub>258</sub> N <sub>72</sub> O <sub>125</sub> P <sub>20</sub> S	6466.2

5

Table 4. ESMS Data for Oligonucleotides 42–53.<sup>a</sup>

Oligonucleotide	Calculated	Formula	Found
42	6575.5	C <sub>221</sub> H <sub>274</sub> N <sub>72</sub> O <sub>122</sub> P <sub>20</sub> S <sub>2</sub>	6576.5
43	6272.2	C <sub>200</sub> H <sub>256</sub> N <sub>72</sub> O <sub>120</sub> P <sub>20</sub> S <sub>2</sub>	6272.3
45	6543.5	C <sub>219</sub> H <sub>269</sub> N <sub>73</sub> O <sub>121</sub> P <sub>20</sub> S <sub>2</sub>	6542.4
46	6366.3	C <sub>209</sub> H <sub>258</sub> N <sub>72</sub> O <sub>119</sub> P <sub>20</sub> S <sub>2</sub>	6366.7
50	6637.6	C <sub>228</sub> H <sub>271</sub> N <sub>73</sub> O <sub>120</sub> P <sub>20</sub> S <sub>2</sub>	6638.0
51	6529.4	C <sub>215</sub> H <sub>259</sub> N <sub>75</sub> O <sub>122</sub> P <sub>20</sub> S <sub>2</sub>	6529.5
52	6556.5	C <sub>219</sub> H <sub>268</sub> N <sub>74</sub> O <sub>121</sub> P <sub>20</sub> S <sub>2</sub>	6555.9
53	6753.7	C <sub>231</sub> H <sub>271</sub> N <sub>73</sub> O <sub>125</sub> P <sub>20</sub> S <sub>2</sub>	6753.1

<sup>a</sup> Oligonucleotide sequence: Tp\*GCATC<sub>5</sub>AG<sub>2</sub>C<sub>2</sub>AC<sub>2</sub>ATp\*\* <SEQ. ID. NO. 10> where p\* and p\*\* are modified phosphorothioate groups (See Schemes 6 and 7).

10

EXAMPLE 12. *Removal of 5'-terminal 3-hydroxy-2,2-bis(ethoxycarbonyl)propyl-1 protecting group.*

Oligonucleotides 16-22 were dissolved in 0.1 M aqueous piperidine (3 mL and 100 mL for 1 and 20 μmol scale, respectively). The reaction mixture was left for 30 min at room temperature. The solvent was evaporated, and the residue was re-dissolved in water (1 to 5 mL). The target oligonucleotides, 24–31 (Tables 1 and 2) were desalted by reversed phase HPLC.

15

EXAMPLE 13. *Oligonucleotide purification by HPLC.*

The oligonucleotides were analyzed, and, for syntheses on 1 to 2 μmol scale, isolated by reverse phase chromatography on a Waters DeltaPak C18 column (15 μm; 300Å; 3.9×300 mm). As buffers A and B, 0.1 M NH<sub>4</sub>OAc and 80% aqueous MeCN were respectively used at a flow rate 1.5 mL min<sup>-1</sup>. Linear gradients from 0 to 60% B in 40 min (Gradient 1) and 0 to 100% in 40 min (Gradient 2) were employed. Retention times for oligonucleotides 8–31 are presented in Table 1. For desalting, the same C18 column was eluted stepwise with 0.1 M NH<sub>4</sub>OAc (10 min), then water (10 min), and, finally, 50% aqueous MeCN (20 min) to give an oligonucleotide as an ammonium salt. For isolation of oligonucleotides synthesized on 20 μmol scale, Waters DeltaPak

20

25

- 5 C18 column (15  $\mu$ m; 300Å; 25×100 mm) was used with buffer systems as described above at the flow rate 15 mL min<sup>-1</sup>.


EXAMPLE 14. *Oligonucleotides 39 and 40.*

10 Referring to scheme 5, oligonucleotide **37** and N-(1-pyrenylmethyl)iodoacetamide, **44**, were added to DMSO. The reaction was kept for 4 h at 37 °C and diluted with water. The product was isolated and desalted by reverse phase HPLC to give **39** as a triethylammonium salt (Table 1).

Oligonucleotide **37** and 5-iodoacetamidofluorescein, **49**, were reacted and the product was isolated and desalted by reverse phase HPLC to give **40** as a triethylammonium salt (Table 1).

15


Scheme 5



5 EXAMPLE 15. *Oligonucleotide 45.*

Referring to scheme 6, oligonucleotide **43** (36 OD, 1.0 mM solution in 200 mM ethyldiisopropylammonium acetate, pH 7.0, 200  $\mu$ L) and N-(1-pyrenylmethyl)iodoacetamide, **44**, (2.0 mg, 25 mM solution in DMSO, 200  $\mu$ L) were added to DMSO (400  $\mu$ L). The reaction was kept for 4 h at 37 °C and diluted to 10 mL with water. The product was isolated and desalted  
 10 by reverse phase HPLC to give **45** (25 OD, 70%) as a triethylammonium salt (Table 1).

Scheme 6



5 EXAMPLE 16. *Oligonucleotide 46.*

Compound **45** (24 OD) was deprotected with conc. aqueous ammonia (3 mL) for 48 h at 55 °C, and the solution was evaporated. The residue was dissolved in 20% aq DMSO, and the product was isolated and desalted by reverse phase HPLC to give **46** (21.5 OD, 90%) as a triethylammonium salt (Table 4).


10

EXAMPLE 17. *Oligonucleotide 50.*

Referring to scheme 7, oligonucleotide **46** (5 OD) and compound **44** (0.5 mg) were reacted in 50 mM ethyldiisopropylammonium acetate buffer (75% aqueous DMSO; pH 7.0; 200 µL) for 8 h at 37 °C as described for **45**. The solution was diluted to 10 mL with water. The product was isolated and desalted by reverse phase HPLC to give **50** (3.5 OD, 70%) as a triethylammonium salt (Table 4).

15

Scheme 7




Oligonucleotide

Hal-R


50

44


51



52



53



## EXAMPLE 18. Oligonucleotide 51.

Oligonucleotide 46 (5 OD) and 4-chloro-7-nitrobenzofurazan, 47, (0.2 mg) were reacted

5 in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 6 h at 37 °C and diluted to 4 mL with water. The product was isolated and desalted by reverse phase HPLC to give **51** (4 OD, 80%) as a triethylammonium salt (Table 4).

EXAMPLE 19. *Oligonucleotide 52.*

10 Oligonucleotide **46** (5 OD) and monobromobimane, **48**, (0.27 mg) were reacted in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 2 h at 37 °C and diluted to 4 mL with water. The product was isolated and desalted by reverse phase HPLC to give **52** (4.5 OD, 90%) as a triethylammonium salt (Table 4).

15 EXAMPLE 20. *Oligonucleotide 53.*

Oligonucleotide **46** (25 OD) and 5-iodoacetamidofluorescein, **49**, (0.52 mg) were reacted in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 10 h at 37 °C and diluted to 4 mL with water. The product was isolated and desalted by reverse phase HPLC to give **53** (19 OD, 75%) as a triethylammonium salt (Table 4).


20 EXAMPLE 21. *Oligonucleotide 56.*

Referring to scheme 8, oligonucleotide **55** (125 OD) and 5-iodoacetamidofluorescein, **49**, (1.5 mg) were reacted in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 10 h at 37 °C and diluted to 4 mL with water. The product was isolated by  
25 reverse phase HPLC, evaporated, and dissolved in concentrated ammonium hydroxide (5 mL). The solution was heated for 48 h at 55 °C and evaporated. The residue was treated with 5% aqueous AcOH (3 mL) for 15 min, evaporated, and treated with 0.1 M aqueous piperidine (3 mL). The product was isolated and desalted by reverse phase HPLC to give **56** (75 OD, 60%) as a triethylammonium salt (Table 5).

30



Scheme 8



5 Table 5. Characterization of oligonucleotides **55–59**.<sup>a</sup>

	ESMS, calculated	Molecular Formula	ESMS, found
<b>55</b>	6886.8	C <sub>231</sub> H <sub>291</sub> N <sub>72</sub> O <sub>131</sub> P <sub>21</sub> S <sub>2</sub>	6885.9
<b>56</b>	6562.2	C <sub>212</sub> H <sub>259</sub> N <sub>72</sub> O <sub>127</sub> P <sub>21</sub> S <sub>2</sub>	6562.9
<b>57</b>	6833.6	C <sub>231</sub> H <sub>272</sub> N <sub>73</sub> O <sub>128</sub> P <sub>21</sub> S <sub>2</sub>	6834.6
<b>58</b>	6752.4	C <sub>222</sub> H <sub>269</sub> N <sub>74</sub> O <sub>129</sub> P <sub>21</sub> S <sub>2</sub>	6753.1
<b>59</b>	6949.6	C <sub>234</sub> H <sub>272</sub> N <sub>73</sub> O <sub>133</sub> P <sub>21</sub> S <sub>2</sub>	6950.8


<sup>a</sup> Oligonucleotide sequence: p\*TCATC<sub>5</sub>AG<sub>2</sub>C<sub>2</sub>AC<sub>2</sub>ATp\*\* <SEQ. ID. NO. 11> where p\* and p\*\* are modified phosphorothioate groups (See Schemes 8 and 9).

10 EXAMPLE 22. *Oligonucleotide 57*.

Referring to scheme 9, oligonucleotide **56** (7 OD) and compound **44** (0.7 mg) were reacted in 50 mM ethyldiisopropylammonium acetate buffer (75% aqueous DMSO; pH 7.0; 250 μL) for 8 h at 37 °C as described for **45**. The solution was diluted to 10 mL with water. The product was isolated and desalted by reverse phase HPLC to give **57** (5.2 OD, 75%) as a

15 triethylammonium salt (Table 5).

Scheme 9



Oligonucleotide	Hal—R
57	44
58	48
59	49

#### EXAMPLE 23. Oligonucleotide 58.

Oligonucleotide **56** (10 OD) and monobromobimane, **48**, (0.53 mg) were reacted in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 2 h at 37 °C and diluted to 4 mL with water. The product was isolated and desalted by reverse phase HPLC to give **58** (8 OD, 80%) as a triethylammonium salt (Table 5).

#### EXAMPLE 24. Oligonucleotide 59.

Oligonucleotide **56** (15 OD) and 5-iodoacetamidofluorescein, **49**, (0.5 mg) were reacted in 0.1 M ethyldiisopropylammonium acetate buffer (60% aq MeCN; pH 7.0; 400  $\mu$ L) for 10 h at 37 °C and diluted to 4 mL with water. The product was isolated and desalted by reverse phase HPLC to give **59** (10.5 OD, 70%) as a triethylammonium salt (Table 5).

5

## EXAMPLE 25

## Synthesis of Dialkylglycerol Linker

10 **1,2-Di-O-hexadecyl-rac-glycerol succinimidyl carbamate (102)** Referring to Scheme 10, 1,2-Di-O-hexadecyl-rac-glycerol (10.00 g, 18.5 mmol) was dissolved in anhydrous  $\text{CH}_2\text{Cl}_2$  (150 ml). To the solution were added disuccinimidyl carbonate 7.11 g, 27.7 mmol),  $\text{Et}_3\text{N}$  (10.0 ml), and MeCN (50 ml). The reaction mixture was stirred at room temperature under Ar for 6.5 h and then evaporated to dryness. The residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (300 ml). It was washed with  
15 saturated  $\text{NaHCO}_3$  aqueous solution ( $3 \times 100$  ml) and with saturated NaCl aqueous solution ( $3 \times 100$  ml). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated to dryness. Compound **102** (12.60 g) was obtained as colorless powder after drying in high vacuum, which was directly used for the next step without further purification.

20 **(1S,2S)-(+)-2-(Fmoc- $\epsilon$ -aminocaproylamido)-1-phenyl-1,3-propanediol (104)** To the solution of Fmoc- $\epsilon$ -aminocaproyl succinimidyl carbamate (**103**, 97% of purity, 27.8 g, 60 mmol) in  $\text{CH}_2\text{Cl}_2$  (300 ml) was added anhydrous pyridine followed by (1S,2S)-(+)-2-amino-1-phenyl-1,3-propanediol (10.08, 60 mmol). The reaction mixture was stirred at room temperature under Ar for 7 h. Meanwhile, white solid was produced. This insoluble material was filtered out. The  
25 filtrate was evaporated to give oil, which was dried in high vacuum overnight furnishing a yellowish powder **104** (31.0 g). For analysis 2.5 g of the crude product was dissolved in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (9:1) and applied to FC (silica gel, column  $3 \times 15$  cm). The column was washed with  $\text{CH}_2\text{Cl}_2$ , 100 ml;  $\text{CH}_2\text{Cl}_2/\text{Me}_2\text{CO}$  9:1, 500 ml. And the compound **104** was eluted with  $\text{CH}_2/\text{MeOH}$  9:1, 200 ml.

30

**(1S,2S)-(+)-2-{ $\epsilon$ -[(1,2-di-O-hexadecyl-rac-glyceroyl)carbonyl]-aminocaproyl}-amido-1-phenyl-1,3-propanediol (105)** The compound **104** (9.04 g, 18 mmol) was dissolved in DMF


5 (108 ml). To this solution piperidine (27 ml) was added. The reaction mixture was stirred at room temperature for 1 h. and then evaporated to dryness. The residue was dissolved in pyridine (36 ml). To this solution was added a solution of 1,2-Di-O-hexadecyl-rac-glycerol succinimidyl carbamate (**102**, 12.92 g, 18 mmol) in  $\text{CH}_2\text{Cl}_2$  (135 ml). The mixture was stirred at room temperature for 5.5 h. and evaporated to dryness. The residue was dried overnight in high vacuum furnishing a yellowish solid. The solid was dissolved in  $\text{CH}_2\text{Cl}_2$  (1000 ml). It was washed with saturated NaCl aqueous solution ( $3 \times 300$  ml). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated to give oil. The residue was applied to FC (silica gel, column  $12 \times 20$  cm),  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (9:1), 2000 ml; yielding compound **105** (7.10 g, 47%) as colorless powder.

15 Compound **105** (6.98 g, 8.2 mmol) was co-evaporated with anhydrous pyridine three times and then dissolved in pyridine (40 ml). To this solution DMTCI (3.34 g, 9.8 mmol) was added under stirring at room temperature in three portions over 7 h. The reaction mixture was stirred at room temperature for another 15 h. The excess DMTCI was decomposed by adding MeOH (20 ml). The solution was poured into saturated  $\text{NaHCO}_3$  aqueous solution (400 ml), shaken and separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 120$  ml). The combined organic layer was washed with saturated NaCl aqueous solution ( $3 \times 200$  ml) and then dried over  $\text{Na}_2\text{SO}_4$ . The solid was filtered out. The filtrate was evaporated to dryness giving a gel, which was applied, to FC (silica gel, column  $12 \times 20$  cm). The column was eluted with  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (95:5 containing drops of Et N) furnishing compound **106** (6.89 g, 73%) as colorless foam.

(1S,2S)-(+)-2-{ $\epsilon$ -[(1,2-di-O-hexadecyl-rac-glyceroxy)carbonyl]-aminocaproyl}-amido-3-[(4,4'-dimethoxytrityl)oxy-1-phenylpropanol 1-O-(2cyanoethyl diisopropylphosphoramidite) (**107**) To the solution of the compound **106** (3.44 g, 3 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (25 ml) were added ( $i\text{Pr}$ ) $_2\text{Net}$  (1.05 ml, 6 mmol) and 2-cyanoethyl-N,N'-diisopropylaminochlorophosphine (0.87 ml, 3.9 mmol) under Ar. The reaction mixture was stirred at room temperature for 1 h. It was then poured into 5%  $\text{NaHCO}_3$  aqueous solution,

- 5 shaken and separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3  $\times$  60 ml). The combined organic layer was washed with 5%  $\text{NaHCO}_3$  aqueous solution (100 ml) and saturated NaCl aqueous solution (2  $\times$  120 ml), dried over  $\text{Na}_2\text{SO}_4$ . The solid was filtered out. The filtrate was evaporated to dryness giving a gel that was further dried in high vacuum furnishing a yellow foam (4.77 g). It was applied to FC (silica gel, column 5  $\times$  15 cm):  $\text{CH}_2\text{Cl}_2/\text{Me}_2\text{CO}$  (9:1), 2000
- 10 ml, yielding **107** (1.0 g) as colorless foam. Also, the starting material **106** (1.0 g) was recovered.

Scheme 10



DSC: Disuccinimidyl carbonate; DEC: 1-(3-Dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride

## 5 EXAMPLE 26

## Synthesis of Cholesterol Linker

**$\epsilon$ -N-Cholesteryloxycarbonylaminocaproic acid (109)** Referring to Scheme 11, the  $\epsilon$ -aminocaproic acid (3.93 g, 30 mmol) was suspended in pyridine (60 ml). The flask was flushed with nitrogen and to the mixture was added N,O-bis(trimethylsilyl)acetamide (10 ml, 70 mmol) under stirring. The reaction mixture was stirred at room temperature for 35 min. and then cooled in ice bath. Cholesteryl chloroformate (13.5 g, 30 mmol) was added into the reaction mixture in two portions over 2 h. The reaction was continued by stirring at room temperature for another 4 h. 2% HCl aqueous solution (150 ml) was added under cooling with ice bath. The mixture was stirred for 5 min. and then poured into a separating funnel. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  150 ml). The combined organic layer was washed with 2% HCl aqueous solution (2  $\times$  120 ml) and with saturated NaCl aqueous solution (2  $\times$  120 ml), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to dryness giving a yellow foam (109, 14.44 g).

20

**(1S,2S)-(+)-2-[ $\epsilon$ -(N-Cholesteryloxycarbonylamino)caproylamino]-1-phenyl-1,3-propanediol (110)** To the solution of compound 109 (crude, 14.44 g) in pyridine (50 ml) 2-(Diethylamino)ethyl chloride hydrochloride (DEC, 6.03 g, 31.5 mmol) was added under nitrogen at room temperature. The mixture was stirred at room temperature under nitrogen for 0.5 h. (1S,2S)-(+)-2-Amino-1-phenyl-1,3-propanediol (5.01 g, 30.0 mmol) was then added followed by another portion of pyridine (10 ml). The reaction mixture was stirred at room temperature for 19 h. It was evaporated to dryness. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2550 ml), washed with saturated NaHCO<sub>3</sub> aqueous solution (2  $\times$  120 ml) followed by saturated NaCl aqueous solution (2  $\times$  150 ml). The combined aqueous layer was re-extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  200 ml). The combined organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give a yellow foam 110 (16.93 g, 81%). For analysis, the crude product (5.67 g) was purified further by FC (silica gel, column 3  $\times$  12 cm): CH<sub>2</sub>Cl<sub>2</sub>, 400 ml; CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95:5), 250 ml; CH<sub>2</sub>Cl<sub>2</sub>/MeOH (9:1), 400 ml furnishing slight yellow powder 110 (4.89 g).

30

5

**(1S,2S)-(+)-3-[(4,4'-Dimethoxytrityl)oxy]-2-[ε-(N-cholesteryloxycarbonylamino)-caproylamino]-1-phenylpropanol (111)** The compound **110** (crude, 12.44 g) was co-evaporated with pyridine (anhydrous, 3 × 20 ml) and dissolved in anhydrous pyridine (50 ml). To the solution DMTCl (5.62 g, 16.6 mmol) was added in three portions over 7 h. The reaction mixture was stirred at room temperature for additional 14 h. The excess DMTCl was then decomposed by adding MeOH (12 ml) and stirring for 10 min. The resulting solution was poured into 5% NaHCO<sub>3</sub> aqueous solution (150 ml). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 × 150 ml). The organic layer was washed by saturated NaCl aqueous solution (200 ml), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated to dryness. The residue was applied to FC (silica gel, column 5 × 8 cm): CH<sub>2</sub>Cl<sub>2</sub>/Me<sub>2</sub>CO (95:5) 750 ml, CH<sub>2</sub>Cl<sub>2</sub>/Me<sub>2</sub>CO (9:1) 750 ml, yielding **111** (11.35 g) as yellowish foam.

20


**(1S,2S)-(+)-3-[(4,4'-Dimethoxytrityl)oxy]-2-[ε-(N-cholesteryloxycarbonylamino)-caproylamino]-1-phenylpropanol 1-O--(2cyanoethyl diisopropylphosphoramidite) (112)** To the solution of the compound **111** (9.95 g, 10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (75 ml) were added (iPr)<sub>2</sub>NEt (3.50 ml, 20 mmol) and 2-cyanoethyl-N,N'-diisopropylaminochlorophosphine (2.90 ml, 13 mmol) under Ar. The reaction mixture was stirred at room temperature for 1 h and was then diluted by adding CH<sub>2</sub>Cl<sub>2</sub> (100 ml). The solution was poured into 5% NaHCO<sub>3</sub> aqueous solution (200 ml), shaken and separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (150 ml). The combined organic layer was washed with 5% NaHCO<sub>3</sub> aqueous solution (2 × 100 ml) and saturated NaCl aqueous solution (3 × 100 ml), dried over Na<sub>2</sub>SO<sub>4</sub>. The solid was filtered out. The filtrate was evaporated to dryness. The residue was applied to FC (silica gel, column 5 × 15 cm): CH<sub>2</sub>Cl<sub>2</sub>/Me<sub>2</sub>CO (95:5), 1000 ml, yielding **112** (6.88 g, 58%) as colorless foam.

25



5

Scheme 11



BSA: Bis(trimethylsilyl)acetamide; DEC: 1-(3-Dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride

10

## EXAMPLE 27

### Synthesis of Solid Support with Immobilized Cholesterol

15

(1S,2S)-(+)-3-[(4,4'-Dimethoxytrityl)oxy]-2-[ε-(N-cholesteryloxycarbonylamino)-caproylamino]-1-phenylpropanyl succinate (113) Referring to Scheme 12, to the solution of the compound 111 (3.98 g, 4 mmol) were added DMAP (0.38 g, 3.2 mmol), succinic anhydride (0.6 g, 6 mmol), and Et<sub>3</sub>N (0.54 ml, 4 mmol). The reaction mixture was stirred at room temperature for 3.5 h. The solution was diluted by CH<sub>2</sub>Cl<sub>2</sub> (150 ml). It was then washed with 10% citric acid aqueous solution (2 × 150 ml) and saturated NaCl aqueous solution (2 × 150 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated to dryness giving a yellow foam (4.33 g).

20

5

**Polymer Support (114)**

(i) CPG Support: To the mixture of the compound **113** (0.5 g, 0.45 mmol), DMAP (0.58 g, 4.5 mmol), and MeCN (2.5 ml) was added the solution of DTNP (0.14 g, 0.45 mmol) in MeCN (1.8 ml) and 1,2-dichloroethane (0.7 ml). After the mixture was stirred at room temperature under nitrogen for 10 min. the solution of TPP (0.12 g, 0.45 mmol) in MeCN (1.2 ml) was added. The stirring was continued for another 10 min. To the mixture resin LCA-CPG (initial loading of NH<sub>2</sub>: 118.9  $\mu$ mol/g, 1.90 g, 0.225 mmol) was added. The mixture was shaken at room temperature for 5 h. and then filtered. The solid residue (resin) was washed alternatively with CH<sub>3</sub>CN (3  $\times$  10 ml), CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  10 ml), and Et<sub>2</sub>O (3  $\times$  10 ml). After drying the resin was capped with Cap A for 30 min followed by washing with CH<sub>2</sub>Cl<sub>2</sub> (2  $\times$  10 ml). It was subsequently capped with Cap B for 30 min. After washing with CH<sub>2</sub>Cl<sub>2</sub> (2  $\times$  10 ml) and Et<sub>2</sub>O (2  $\times$  10 ml), the resin was dried in high vacuum at room temperature overnight yielding polymer support. Loading of the cholesterol (35.2  $\mu$ mol/g) was observed by determination of released DMT absorption at 498 nm.


20

(ii) Prime Support: The compound **113** (1.28 g, 1.17 mmol) and HATU (0.45 g, 1.17 mmol) were dissolved in anhydrous DMF (18 ml). To the solution (iPr)<sub>2</sub>Net (0.53 ml, 3.13 mmol) was added. The mixture was shaken for 5 min and the resin (4g, Primer Support 30, HL, Aminoderivatised Version, Pharmacia Biotech, original loading 146  $\mu$ mol/g) was added. The reaction mixture was shaken at room temperature for 40 h. It was filtered and the residue was washed with DMF (4  $\times$  20 ml) followed by CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  20 ml), dried in high vacuum for 2.5 h. The unreacted amino function was capped with Cap A for 30 min. The resin was washed with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  20 ml). It was then capped with Cap B for 30 min. followed by washing with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  20 ml). The residue was dried in high vacuum at room temperature for 24 h. yielding the polymer support (4.44 g, loading of cholesterol 51.4  $\mu$ mol/g).

30

5

Scheme 12



10

## EXAMPLE 28

15 **Phosphoramidite derived from dihexadecylglycerol containing non-nucleosidic linker**


Referring to Scheme 13, compound 115 (6.98 g, 8.2 mmol) was co-evaporated with anhydrous pyridine three times and then dissolved in pyridine (40 ml). To this solution DMTCI (3.34 g, 9.8 mmol) was added under stirring at room temperature in three portions over 7 h. The reaction mixture was stirred at room temperature for another 15 h. The excess DMTCI was decomposed by adding MeOH (20 ml). The solution was poured into a saturated  $\text{NaHCO}_3$  aqueous solution (400 ml), shaken and separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3  $\times$  120 ml). The combined organic layer was washed with a saturated NaCl aqueous solution (3  $\times$  200 ml) and then dried over  $\text{Na}_2\text{SO}_4$ . The solid was filtered out. The filtrate was evaporated to dryness giving a gel, which was applied, to a silica gel, column 12  $\times$  20 cm. The column was eluted with  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (95:5 containing drops of  $\text{Et}_3\text{N}$ ) furnishing compound 116 (6.89 g,

25

5 73%) as colorless foam.

**Compound (117)** To the solution of the compound **116** (3.44 g, 3 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (25 ml) were added  $(i\text{Pr})_2\text{Net}$  (1.05 ml, 6 mmol) and 2-cyanoethyl- $\text{N},\text{N}'$ -diisopropylaminochlorophosphine (0.87 ml, 3.9 mmol) under Ar. The reaction mixture was  
10 stirred at room temperature for 1 h. It was then poured into 5%  $\text{NaHCO}_3$  aqueous solution, shaken and separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 60$  ml). The combined organic layer was washed with 5%  $\text{NaHCO}_3$  aqueous solution (100 ml) and saturated  $\text{NaCl}$  aqueous solution ( $2 \times 120$  ml), dried over  $\text{Na}_2\text{SO}_4$ . The solid was filtered out. The filtrate was  
15 evaporated to dryness giving a gel that was further dried in high vacuum furnishing a yellow foam (4.77 g). It was applied to FC (silica gel, column  $5 \times 15$  cm):  $\text{CH}_2\text{Cl}_2/\text{Me}_2\text{CO}$  (9:1), 2000 ml, yielding **117** (1.0 g) as colorless foam. Also, the starting material **116** (1.0 g) was recovered.

Scheme 13




DSC: Disuccinimidyl carbonate


## 5           EXAMPLE 29

          The cholesterol- and dialkylglycerol-conjugated oligonucleotides, compounds **120-129**, presented in the Tables 6-9 below were prepared using the phosphoramidites and solid support described above in connection with Examples 27 and 28. The RP HPLC profiles of compound **120** in its crude form prior to removal of the DMT group, in its crude form after removal of the DMT group, and in its final form are shown in FIGS. 1-3, respectively. The RP HPLC profiles of compound **129** in its crude form prior to removal of the DMT group, in its crude form after removal of the DMT group, and in its final form are shown in FIGS. 4-6, respectively. The RP HPLC profiles of compound **126** in its crude form after removal of the DMT group and in its final form are shown in FIGS. 7 and 8, respectively. The RP HPLC profiles of compound **121** in its crude form after removal of the DMT group and in its final form are shown in FIGS. 9 and 10, respectively.


Compound 120



Compound 129



Compound 126





Compound 121




Table 6: Cholesterol-/Dialkylglycerol-Conjugated Oligonucleotides (1)

Cmpd.	Sequence (5' - 3')	Backbone	Quantity (ODs at 260 nm)	Molecular Weight		More Information
				calc.	found	
120	Chol-5'-d(GCC CAA GCT GGC ATC CGT CA) <SEQ. ID. NO. 12>	P=S	58 mg (880 OD)	7139	7137	ICAM-1
121	Diag-5'-d(GCC CAA GCT GGC ATC CGT CA) <SEQ. ID. NO. 12>	P=S	46 mg (816 OD)	7293	7290	ICAM-1

Note: (1) Chol: Cholesterol and Linker, see structure. (2) Diag: Dialkylglycerol and Linker.







Table 7: Cholesterol-Conjugated Oligonucleotides (2)

Cmpd.	Sequence (5' – 3')	Backbone	Quantity (ODs at 260 nm)	Molecular Weight		More Information
				calc.	found	
122	Chol-5'-d(tcc GTC ATC GCT cct cag gg) <SEQ. ID. NO. 13>	P=S	25 mg (456 OD)	8006	8006	Ras-Ha
123	Chol-5'-d(gtc caC CAT TAG CAC gcg gg) <SEQ. ID. NO. 14>	P=S	42 mg (727 OD)	7962	7961	TGF- $\beta$
124	Chol-5'-d(gtc caC* C*AT TAG C*AC* gcg gg) <SEQ. ID. NO. 15>	P=S	42 mg (789 OD)	8018	8018	TGF- $\beta$
<p>Note: (1) Chol: Cholesterol and Linker, see structure.</p> <p>(2) t: 2'-O-MOE-rT; c: 2'-O-MOE-5-Me-rC; a: 2'-O-MOE-rA; g: 2'-O-MOE-rG; C*: 5-Me-dC</p>						



5 Table 8: Cholesterol-Conjugated Oligonucleotides (3)

Cmpd.	Sequence (5' - 3')	Backbone	Quantity (ODs at 260 nm)	Molecular Weight		More Information
				calc.	found	
125	Chol-5'-(cct ctt acc tea gtt aca) <SEQ. ID. NO. 16>	P=S	67 OD	7845	7846	RNA-splicing
126	Chol-5'-(gct att acc tta acc cag) <SEQ. ID. NO. 17>	P=S	70 OD	7880	7881	RNA-splicing
<p>Note: (1) Chol: Cholesterol and Linker, see structure.            (2) t: 2'-O-MOE-rT; c: 2'-O-MOE-5-Me-rC;            a: 2'-O-MOE-rA; g: 2'-O-MOE-rG.</p> <p>Kole705: 5'-(CCU CUU ACC UCA GUU ACA) &lt;SEQ. ID. NO. 20&gt;            Kole654: 5'-(GCU AUU ACC UUA ACC CAG) &lt;SEQ. ID. NO. 21&gt;</p> <div style="text-align: center;">  <p>Cholesterol and Linker</p> </div>						

Docket No.: ISIS-4723

PATENT

Table 9: Cholesterol-Conjugated Oligonucleotides (4)

Cmpd.	Sequence (5' - 3')	Backbone	Quantity	MW		More Information
				calc.	Found	
127	Chol-5'-d(ctg ctA GC*C* TC*T GGA tt ga) <SEQ. ID. NO. 18>	P=S	82 mg	7986	7993	PTEN
128	Chol-5'-d(ctt ctG GC*A TC*C* GGT tta ga) <SEQ. ID. NO. 19>	P=S	84 mg	7986	7994	PTEN

5

Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of the invention. It is therefore intended that the appended claims cover all equivalent variations as fall within the true scope and spirit of the invention.

10